arXiv:gr-qc/9405046v1 22 May 1994

Short title: **Newtonian Gravitation as a Gauge Theory.**

P.A.C.S. number:+04.50,04.90,03.50

STANDARD AND GENERALIZED NEWTONIAN GRAVITIES AS "GAUGE" THEORIES OF THE EXTENDED GALILEI GROUP - I: THE STANDARD THEORY

R. DE PIETRI[1]

Department of Physics and Astronomy, University of Pittsburgh,
Pittsburgh, PA 15260, USA

L. LUSANNA

I.N.F.N., Sezione di Firenze
Largo E. Fermi 2, 50127 Arcetri (FI), Italy

and

M. PAURI[1]

Center for Philosophy of Science - University of Pittsburgh
817D Cathedral of Learning, Pittsburgh, PA 15260, USA

Abstract

Newton's standard theory of gravitation is reformulated as a *gauge* theory of the *extended* Galilei Group. The Action principle is obtained by matching the *gauge* technique and a suitable limiting procedure from the ADM-De Witt action of general relativity coupled to a relativistic mass-point.

---

[1] On leave from: Dipartimento di Fisica - Sezione di Fisica Teorica, Università di Parma, 43100 Parma, Italy, and I.N.F.N., Sezione di Milano, Gruppo Collegato di Parma

1 Introduction

The present work is the first of a series in which a suitable reformulation of the *gauge* procedure is exploited for dealing with classical non-relativistic systems. In particular, this is the first of two papers in which the *gauge* technique is applied to *extended* Galilei group. The general scope of this treatment is to reformulate firstly standard Newton‘s theory as a *general manifestly-covariant* Galilean *gauge* theory, and, secondly, to seek possible generalizations of it.

As is well-known, a geometrical four-dimensional formulation of Newton's gravitational theory has been developed already in the thirties by Elie Cartan [1]. More recent formulations of the classical theory of gravitation in geometrical terms have been proposed by Havas [2], Anderson [3], Trautman [4], Künzle [5] and Kuchař [6]. Analyses of the classical theory as a non-relativistic limit of general relativity has been made by Dautcourt [7], Künzle [5], Ehlers [8], Malament [9] and others. In all these papers, the Newtonian theory is reobtained by describing its inertial-gravitational structure in terms of an affine connection compatible with the temporal flow t_μ and a rank-three spatial metric $h^{\mu\nu}$. While the curvature of the four-dimensional affine connection is different from zero because of the presence of matter, the Newtonian flatness of the absolute three-space is guaranteed by the further requirement that Poisson's equation be satisfied, in the covariant form $R_{\mu\nu} = 4\pi G\rho(z)t_\mu t_\nu$, where $R_{\mu\nu}$ is the Ricci tensor of the affine connection and $\rho(z)$ is the matter density. In this way the four-dimensional description is *dynamical*, while the three-dimensional one is not. A further typical feature of this geometrical formulation is the fact that, unlike the case of general relativity in which there is a unique compatible affine connection, the curved four-dimensional affine structure can be separated out in a flat affine (*inertial*) structure and a gravitational (*force*) potential; this splitting, however, cannot be done in a unique way, unless special boundary conditions are externally provided.

While the four-dimensional point of view about Newton's theory shows remarkable geometrical insights and even provides a better foundation for Newtonian kinematics than does the traditional point of view (see, in this connection Earman and Friedman [10]), it does not lend itself to any easy generalization. And since what we want to obtain in the end is precisely a true generalization of Newton's theory allowing for a dynamical three-space, we will adopt here a completely independent procedure based on a three-dimensional level of description from the beginning. With this in view, the present paper should be read also as a first necessary step towards the searched generalized formulation. Our approach develops through the following steps:

(1) First, we exploit the *gauge* methodology originally applied by Utiyama [11] to the Lorentz group within the field theoretic framework, in order to find all the inertial-gravitational fields which can be coupled to a non-relativistic mass-point. To achieve this result, we apply a suitably adapted Utiyama procedure to the Galilei group. Specifically, we consider a Galilei invariant Action corresponding to the *projective* canonical realization which describes a free particle of mass m. The requirement of invariance (properly *quasi*-invariance, according to what usually obtains in the case of groups with non trivial cohomology structure) of the Action under *localized* Galilei transformations, leads directly to the following results: (a) the introduction of eleven compensating *gauge* fields (one more than the order of the standard Galilei group due to the *central extension* of it); (b) the characterization of their Galilean transformation properties, and ; (c) the explicit form of the Action describing the dynamics of the mass-point interacting with the *gauge* fields playing the role of external fields. A geometrical interpretation of these latter fields is then exhibited by evidentiating their relation to the so-called *Galilei* and *Newtonian* Structures studied by Künzle [5] and Kuchař [6].

(2) Second, we look for a possible *field* Action capable of describing the *dynamics* of these fields. In realizing this program we are guided by the following facts: (a) The non-relativistic limit of the relativistic Lagrangian for a mass-point in a pseudo-Riemannian space is precisely the Galilei matter Lagrangian we have already obtained through the *gauge* procedure; (b) the non-relativistic limit of Einstein equations leads to the geometric Cartan structure with Newton's equations; on the other hand: (c) none among the existing four-dimensional formulations of Newtonian gravitation is cast in a variational form.

The explicit construction of the fundamental Galilei Action is performed by matching the results obtained through the above *gauge* technique and a suitable non-relativistic limiting procedure (for $c^2 \to \infty$) from the four-dimensional level. Precisely, the limiting procedure is applied to the Einstein-Hilbert-De Witt action for the gravitational field plus a matter action corresponding to a single mass point, under the assumption of the existence of a global 3+1 splitting of the total Action, and of a suitable parametrization of the 4-metric tensor in terms of powers of c^2. Once the expansion in powers of $1/c^2$ is explicitly calculated, we make the *Ansatz* of identifying the wanted Galilean Action A with the $zero^{th}$ order term of the expansion itself.

The resulting Action contains 27 fields, i.e., 16 fields *over and above* the *gauge* fields obtained through the *gauge* technique. These fields are not coupled to matter and lack *a-priori* any definite transformation property. Their role is nonetheless essential (in the spirit of the Einstein-Kretschmann debate[12], one would say) as *auxiliary* fields, to the effect that they guarantee a *Galilean general-covariance* of the three-dimensional theory (where of course the *absolute* nature of *time* is preserved). In fact, once appropriate transformation properties for the *auxiliary* fields are postulated, the fundamental Galilei Action turns out to be *quasi-invariant* under the *local* Galilei transformations.

As expected, a constraint analysis shows that the theory has *no physical field degrees of freedom* so that it is essentially Newton's theory expressed as a *gauge* invariant theory or, stated in other words, as a theory cast in a form valid in arbitrary (*absolute time* respecting) *Galilean reference frames*. This formulation implies, of course, flatness of the three-space metric g_{ij} (expressed through the vanishing of the three-dimensional Ricci tensor: $R_{ij} = 0$) and validity of the Poisson equation in a suitable Galilei-covariant form (see Section 7.1). As far as we know, a manifestly-covariant formulation of Newton's theory of gravitation has never been proposed until now.

In Section 2, the free mass-point realization of the extended Galilei group is expounded together some preliminaries and notations. Section 3 is devoted to the *gaugeization* of the group: *gauge* compensating fields and their group transformation properties are derived by imposing *quasi-invariance* of the action. The equations of motion of the mass-pont in presence of the *gauge* "external" fields are discussed in Section 4 together with a proper characterization of the relations between various kinds of "observers" and "forms" of the fields. Section 5 is dedicated to a summary of the known facts about Galilean and Newtonian geometrical structures and to the correspondence between these latter and our *gauge* fields. In section 6 the main *Ansatz* for the selection of the fundamental Galilean Action is discussed. Section 7 is devoted to a discussion of the constraint analysis of the covariant form of Newton's theory in *arbitrary - absolute time respecting* reference systems (Section 7.1) and in *Galilean* reference systems (Section 7.2), respectively.

2 Preliminaries on the Galilei free mass-point realization

The Lagrangian and the action for a free non relativistic mass-point can be written,

$$L_M = \frac{1}{2} m \delta_{ij} \dot{x}^i \dot{x}^j \ , \ \ \mathcal{A}_M = \int_{t_1}^{t_2} dt \ \ L_M \ , \left(\delta_{ij} = \left\{ \begin{array}{ll} 1, & i=j \\ 0, & i \neq j \end{array} \right. \right) \ , \tag{2.1}$$

respectively.

The variational principle $\delta \mathcal{A}_M = 0$, with variations which vanish at the end points, gives the Euler-Lagrange equations:

$$\mathcal{EL}_i \equiv \frac{\partial L_M}{\partial x^i} - \frac{d}{dt} \frac{\partial L_M}{\partial \dot{x}^i} = -m \delta_{ij} \ddot{x}^j \overset{\circ}{=} 0 \ , \tag{2.2}$$

where $\overset{\circ}{=}$ means that the equality is satisfied on the extremals.

The infinitesimal Galilei transformations of the configuration variables will be written as:

$$\left\{ \begin{array}{ll} \delta t & = -\varepsilon \\ \delta x^i & = \varepsilon^i + c_{jk}{}^i \omega^j x^k - v^i t \\ \delta \dot{x}^i & = c_{jk}{}^i \omega^j \dot{x}^k - v^i \ \ , \end{array} \right. \tag{2.3}$$

where $\varepsilon, \varepsilon^i, \omega^i, v^i$ are the infinitesimal parameters of time translation, space translations, space rotations and pure Galilei transformations (Galilei boosts), respectively, and the $c_{jk}{}^i$'s are the standard structure-constants of the SO(3) group.

We remark that, for a given infinitesimal time transformation $t \to t^* = t + \delta t$, $f(t) \to f^*(t^*)$, the symbol δ means $\delta f(t) \equiv f^*(t^*) - f(t)$. On the other hand, the *equal time* infinitesimal transformation will be denoted by $\delta_0 f(t) \equiv f^*(t) - f(t) = \delta f - \delta t \cdot \frac{df}{dt}$. The *equal-time* configuration variables transformations corresponding to the transformation (2.3) are:

$$\begin{cases} \delta_0\, t &= 0 \\ \delta_0 x^i &= \varepsilon^i + c_{jk}{}^i \omega^j x^k - t v^i + \varepsilon \dot{x}^i \\ \delta_0 \dot{x}^i &= c_{jk}{}^i \omega^j \dot{x}^k - v^i + \varepsilon \ddot{x}^i \quad . \end{cases} \tag{2.4}$$

Note that, while δ_0 commutes with time derivative, so that

$$\delta_0 \frac{df(t)}{dt} = \frac{d}{dt}\delta_0 f(t) \quad , \tag{2.5}$$

we have instead

$$\begin{aligned} \delta \frac{df(t)}{dt} &= \delta_0 \frac{df(t)}{dt} + \frac{d^2 f(t)}{dt^2}\delta t \\ &= \frac{d}{dt}\delta f(t) - \frac{df(t)}{dt}\frac{d(\delta t)}{dt} \quad . \end{aligned} \tag{2.6}$$

Finally, notice that, since the Galilei transformations are not, in general, fixed-time transformations, the variation of the action must be explicitly written in the form:

$$\Delta \mathcal{S}_M = \int_{t_1}^{t_2} dt \left[\delta L_M + L_M \frac{d}{dt}\delta t\right] = \int_{t_1}^{t_2} dt \left[\delta_0 L_M + \frac{d}{dt}(L_M \delta t)\right] \quad . \tag{2.7}$$

Under the transformations (2.3), we have:

$$\Delta \mathcal{S}_M = \int_{t_1}^{t_2} dt \left[\frac{d}{dt}\left(-m\delta_{ij} x^i v^j\right)\right] \quad , \tag{2.8}$$

so that the action (2.1) results *quasi-invariant* under them. As a consequence of this *quasi-invariance* of the action, we have the Noether identity:

$$\dot{G} \equiv -\delta_0 x^i \ \mathcal{EL}_i \overset{\circ}{=} 0 \quad , \tag{2.9}$$

where the constant of the motion G is given by:

$$G = \frac{\partial L_M}{\partial \dot{x}^i}\delta_0 x^i - \varepsilon L_M + m\delta_{ij} x^i v^j \ . \tag{2.10}$$

Before proceeding, let us fix some notation in connection with more general classes of functions that we will have to consider; precisely: 1) $f(\boldsymbol{z}, t)$, with $\boldsymbol{z}$ and t independent variables; and 2) $f(\boldsymbol{x}(t), t)$.

The variations in case 1) when $t \to t^* = t + \delta t$, $\boldsymbol{z} \to \boldsymbol{z}^* = \boldsymbol{z} + \delta \boldsymbol{z}$, will be denoted by

$$\begin{aligned} \delta f(\boldsymbol{z}, t) &= f^*(\boldsymbol{z}^*, t^*) - f(\boldsymbol{z}, t) \\ \delta_0 f(\boldsymbol{z}, t) &= f^*(\boldsymbol{z}, t) - f(\boldsymbol{z}, t) = \\ &= \delta f(\boldsymbol{z}, t) - \frac{\partial f(\boldsymbol{z}, t)}{\partial z^k} \delta z^k - \frac{\partial f(\boldsymbol{z}, t)}{\partial t} \delta t \quad . \end{aligned} \tag{2.11}$$

On the other hand, in case 2), it is convenient to distinguish three kinds of variations: if

$$\begin{cases} t & \to \quad t^* = t + \delta t \\ \boldsymbol{x}(t) & \to \quad \boldsymbol{x}^*(t^*) = \boldsymbol{x}(t) + \delta \boldsymbol{x}(t) = \boldsymbol{x}(t) + \delta_0 \boldsymbol{x}(t) + \dfrac{d\boldsymbol{x}(t)}{dt} \delta t \quad , \end{cases} \tag{2.12}$$

we shall define:

$$\begin{aligned} \delta f(\boldsymbol{x}(t), t) &= f^*(\boldsymbol{x}^*(t^*), t^*) - f(\boldsymbol{x}(t), t) \\ \delta_0 f(\boldsymbol{x}(t), t) &= f^*(\boldsymbol{x}(t), t) - f(\boldsymbol{x}(t), t) = \\ &= \delta f(\boldsymbol{x}(t), t) - \frac{\partial f(\boldsymbol{x}(t), t)}{\partial x^k} \delta x^k(t) - \frac{\partial f(\boldsymbol{x}(t), t)}{\partial t} \delta t \\ \delta_{0[t]} f(\boldsymbol{x}(t), t) &= f^*(\boldsymbol{x}^*(t), t) - f(\boldsymbol{x}(t), t) = \\ &= \delta f(\boldsymbol{x}(t), t) - \left[\frac{\partial f(\boldsymbol{x}(t), t)}{\partial x^k} \frac{dx^k(t)}{dt} + \frac{\partial f(\boldsymbol{x}(t), t)}{\partial t} \right] \delta t \end{aligned} \tag{2.13}$$

Let us now turn to the Hamiltonian formalism. The canonical momenta and the Hamiltonian function are ($\bar{f}(p, q)$ denotes a function in phase-space),

$$\begin{aligned} p_i &= \frac{\partial L_M}{\partial \dot{x}^i} = m \, \delta_{ij} \dot{x}^j \\ \bar{H} &= p_i \dot{x}^i - L_M = \tfrac{1}{2m} \, \delta^{ij} p_i p_j \quad , \end{aligned} \tag{2.14}$$

respectively. Then, the conserved quantity G becomes:

$$\begin{aligned} \bar{G} &= \varepsilon \bar{H} + \varepsilon^i p_i + \omega^i \bar{J}_i + v^i \bar{K}_i \\ &= \varepsilon \bar{H} + (\eta^i - t v^i) p_i + m \delta_{ij} v^i x^j \overset{\circ}{=} 0 \quad , \end{aligned} \tag{2.15}$$

where, for future convenience, we have introduced the infinitesimal transformation descriptors

$$\eta^i(x^j) = \varepsilon^i + c_{jk}{}^i \omega^j x^k \quad . \tag{2.16}$$

From Eq.(2.15), we obtain the following independent constants of the motion:

$$\bar{H} \ , \quad p_i \ , \quad \bar{K}_i = m \delta_{ij} \ x^j - t \ p_i \ , \quad \bar{J}_i = c_{ij}{}^k x^j p_k \ . \tag{2.17}$$

The constants of motion (2.17) provide a *projective* canonical realization of the Lie algebra of the Galilei group in terms of Poisson brackets:

$$\begin{array}{ll} \{\mathcal{K}_i, \mathcal{H}\} & = \mathcal{P}_i \\ \{\mathcal{K}_i, \mathcal{P}_j\} & = \delta_{ij} m \\ \{\mathcal{P}_i, \mathcal{J}_j\} & = c_{ij}{}^k \mathcal{P}_k \\ \{\mathcal{K}_i, \mathcal{J}_j\} & = c_{ij}{}^k \mathcal{K}_k \\ \{\mathcal{J}_i, \mathcal{J}_j\} & = c_{ij}{}^k \mathcal{J}_k \ , \end{array} \tag{2.18}$$

where:

$$\mathcal{H} = \bar{H} \ , \quad \mathcal{P}_i = p_i \ , \quad \mathcal{K}_i = \bar{K}_i \ , \quad \mathcal{J}_i = \bar{J}_i \ . \tag{2.19}$$

Alternatively, the realization can be considered as a *true* realization [13] of the centrally-extended Galilei group via the *central charge* $\mathcal{M} = m$.

The generator of the *equal-time* Galilei transformation in phase-space, say $\bar{\delta}_0$, is given by the expression $\bar{G}$ defined in eq.(2.15). We have

$$\left\{ \begin{array}{lcl} \bar{\delta}_0 t & = & 0 \\ \bar{\delta}_0 x^i & = & \{x^i, \bar{G}\} = \varepsilon \dfrac{1}{m} \delta^{ij} p_j + \eta^i \\ & & \Rightarrow \ \ \bar{\delta}_0 x^i \Big|_{p=\partial L_M/\partial \dot{x}} = \delta_0 x^i = \delta x^i + \varepsilon \dot{x}^i \quad . \end{array} \right. \tag{2.20}$$

Notice that, within the Hamiltonian formalism, the variation $\delta_0 \dot{x}^i$ can be obtained only by using the equations of motion. In fact, we have:

$$\begin{array}{ll} \delta_0 \dot{x}^i & = \dfrac{\partial \eta^i(x)}{\partial x^k} \dot{x}^k - v^i - \varepsilon \ddot{x}^i = \dfrac{\partial \eta^i(x)}{\partial x^k} \dot{x}^k - v^i - \varepsilon \dfrac{1}{m} \delta^{ij} \mathcal{EL}_j \\ & \overset{\circ}{=} \dfrac{\delta^{ij}}{m} \bar{\delta} p_j \Big|_{p=\partial L_M/\partial \dot{x}} \quad . \end{array} \tag{2.21}$$

Under the transformations (2.4), which are the configuration-space analogues of the transformations (2.20), we have:

$$\Delta \mathcal{S}_M = \int_{t_1}^{t_2} dt \left[\frac{d}{dt} \left(\varepsilon L_M - m \delta_{ij} x^i v^j \right) \right] \ . \tag{2.22}$$

The resulting Noether's constants of the motion are clearly the same as those associated with the transformations (2.3) and (2.20).

This complication can be easily avoided by turning to a re-parameterization invariant formulation of the free mass-point system. Using coordinates $t(\lambda), x^i(\lambda)$, the Lagrangian and the Action become

$$\begin{aligned}\hat{L}_M(\lambda) &= \frac{1}{2}m\frac{\delta_{ij}x'^i(\lambda)x'^j(\lambda)}{t'(\lambda)} \quad ,\\ \hat{\mathcal{S}}_M &= \int_{\lambda_1}^{\lambda_2} d\lambda \;\; \hat{L}_M(\lambda) \quad ,\end{aligned} \tag{2.23}$$

respectively, where $f'(\lambda) \equiv \frac{d}{d\lambda}f(\lambda)$. As before, in this enlarged space, we can define again an infinitesimal transformation $\hat{\delta}$ and an "equal-λ" one, say $\hat{\delta}_0$. The associated Euler-Lagrange equations and canonical momenta are

$$\begin{cases}\widehat{\mathcal{EL}}_t &= \dfrac{d}{d\lambda}\left[\dfrac{m}{2}\dfrac{\delta_{ij}x'^i(\lambda)x'^j(\lambda)}{t'^2(\lambda)}\right] \overset{\circ}{=} 0\\ \widehat{\mathcal{EL}}_i &= -\dfrac{d}{d\lambda}\left[m\dfrac{\delta_{ij}x'^j(\lambda)}{t'(\lambda)}\right] \overset{\circ}{=} 0 \quad ,\end{cases} \tag{2.24}$$

$$\begin{cases}\hat{E} &= -\dfrac{\partial \hat{L}}{\partial t'} = \dfrac{m}{2}\dfrac{\delta_{ij}x'^i(\lambda)x'^j(\lambda)}{t'^2(\lambda)} = \dfrac{1}{2m}\delta^{ij}\hat{p}_i\hat{p}_j\\ \hat{p}_i &= \dfrac{\partial \hat{L}}{\partial x'^i} = m\dfrac{\delta_{ij}x'^j(\lambda)}{t'(\lambda)} = m\delta_{ij}\dot{x}^j(t) = p_i \quad ,\end{cases} \tag{2.25}$$

respectively, where we have defined the Poisson brackets so that:

$$\begin{aligned}\{t(\lambda), \hat{E}(\lambda)\}' &= -1\\ \{x^i(\lambda), \hat{p}_j(\lambda)\}' &= \delta^i_j \quad .\end{aligned} \tag{2.26}$$

In the enlarged phase-space, coordinatized by $(t,\; x^i,\; \hat{E},\; \hat{p}_i)$, we obtain a vanishing canonical Hamiltonian and the first-class constraint

$$\hat{\chi} \equiv \hat{E} - \frac{1}{2m}\delta^{ij}\hat{p}_i\hat{p}_j \approx 0 \quad . \tag{2.27}$$

The constraint $\hat{\chi}$ generates the following transformation of the configurational variables: $\hat{\hat{\delta}}_0 t(\lambda) = -\alpha(\lambda)$, $\hat{\hat{\delta}}_0 x^i(\lambda) = -\alpha(\lambda)\frac{1}{m}\delta^{ij}\hat{p}_j$. This is the re-parameterization gauge transformation $\lambda \rightarrow \lambda - \alpha(\lambda)/t'$. The Lagrangian is obviously *quasi-invariant* under this operation since:

$$\hat{\delta}_0\hat{L}_M = \frac{d}{d\lambda}\left[\frac{-\alpha(\lambda)}{t'}\hat{L}_M\right] \quad . \tag{2.28}$$

The canonical generators of the extended Galilei algebra are now:

$$\hat{\mathcal{H}} = \hat{E} \;,\;\; \hat{\mathcal{P}}_i = \hat{p}_i \;,\;\; \hat{\mathcal{K}}_i = m\;\delta_{ij}\;x^j - t\;\hat{p}_i \;,\;\; \hat{\mathcal{J}}_i = c_{ij}{}^k x^j \hat{p}_k \;,\;\; \hat{\mathcal{M}} = m \;. \tag{2.29}$$

and satisfy the Lie-algebra (2.18) with the primed Poisson-brackets (2.26). Consequently, the generator of the complete phase-space Galilei transformation $\hat{\bar{\delta}}_0$, which is now given by:

$$\begin{aligned} \hat{\bar{G}} &= \varepsilon\hat{E} + \varepsilon^i\hat{p}_i + \omega^i\hat{\bar{J}}_i + v^i\hat{\bar{K}}_i \\ &= \varepsilon\hat{E} + (\eta^i - tv^i)\hat{p}_i + m\delta_{ij}v^ix^j \quad , \end{aligned} \tag{2.30}$$

yields the following "equal-λ" infinitesimal transformations:

$$\begin{cases} \hat{\bar{\delta}}_0\lambda &= 0 \\ \hat{\bar{\delta}}_0 t &= \{t\ ,\hat{\bar{G}}\}' = -\varepsilon = \hat{\delta}_0 t = \delta t \\ \hat{\bar{\delta}}_0 x^i &= \{x^i,\hat{\bar{G}}\}' = \varepsilon^i + c_{jk}{}^i\omega^jx^k - tv^i = \hat{\delta}_0x^i = \delta x^i \quad , \end{cases} \tag{2.31}$$

which coincide with the transformations (2.3).
We have now $\hat{\bar{\delta}}_0\hat{p}_i\Big|_{\hat{p}=\partial\hat{L}_M/\partial x'} = \hat{\delta}_0\left[\frac{m\delta_{ij}x'^j}{t'}\right]$, without any use of Euler-Lagrange equations. On the other hand, under the transformations (2.31), it follows:

$$\hat{\delta}_0\hat{L}_M = \frac{d}{d\lambda}\left[-m\delta_{ij}v^ix^j\right] \quad , \tag{2.32}$$

and

$$\hat{\bar{\delta}}_0\hat{\chi} = 0 \quad , \tag{2.33}$$

so that the canonical generators (2.29) are again constants of the motion. Furthermore, the first-class constraint is Galilei invariant, and the *quasi-invariance* of the Lagrangian is an effect of the *central-charge* term alone.

3 "Gauging" the extended Galilei algebra for the free mass-point

We proceed now to *gauging* the Galilei transformations along the standard line of Utiyama [11]. Since the Newtonian time is absolute, the most general finite transformation allowed for the time coordinate is of the form $t \to t' = t + f(t)$. Therefore a first guess of how to *gauge* the group amounts to replace the complete infinitesimal generators (2.15) and (2.30) with:

$$\begin{aligned} \bar{G} &= \varepsilon(t)\ \bar{H} + \varepsilon^i(\boldsymbol{x},t)\ p_i + \omega^i(\boldsymbol{x},t)\ \bar{J}_i + v^i(\boldsymbol{x},t)\ \bar{K}_i \\ &= \varepsilon(t)\ \bar{H} + [\eta^i(\boldsymbol{x},t)\ - tv^i(\boldsymbol{x},t)\ ]p_i + m\ \delta_{ij}v^i(\boldsymbol{x},t)\ x^j \quad , \end{aligned} \tag{3.1}$$

and

$$\begin{aligned}\hat{\bar{G}} &= \varepsilon(t)\ \hat{E} + \varepsilon^i(\boldsymbol{x},t)\ \hat{p}_i + \omega^i(\boldsymbol{x},t)\ \hat{\bar{J}}_i + v^i(\boldsymbol{x},t)\ \hat{\bar{K}}_i \\ &= \varepsilon(t)\ \hat{E} + [\eta^i(\boldsymbol{x},t)\ - tv^i(\boldsymbol{x},t)]\ \hat{p}_i + m\ \delta_{ij} v^i(\boldsymbol{x},t)\ x^j \quad ,\end{aligned} \tag{3.2}$$

respectively, with ε independent of $\boldsymbol{x}$, and

$$\eta^i(\boldsymbol{x},t) = \varepsilon^i(\boldsymbol{x},t)\ + c_{jk}{}^i \omega^j(\boldsymbol{x},t)\ x^k \quad . \tag{3.3}$$

Notice that we have not absorbed the term $tv^i(\boldsymbol{x},t)$ into $\eta^i(\boldsymbol{x},t)$, as it would be natural from the point of view of the configuration space. Actually, this would not be as much natural in phase-space and, in addition, from the group-theoretical point of view, it would be confusing: it would mix the role of the *central charge* with that of the three-dimensional Euclidean subalgebra.

We see from Eq.(3.1) and Eq.(3.2) that *gauging* the extended Galilei algebra in the case of the mass-point realization is equivalent to *gauging* the algebra generated by energy (respectively *time-translation*, within the re-parameterization invariant picture), linear momentum, and the *central-charge* $\mathcal{M} = m$, with parameters $\varepsilon(t)$, $\eta^i(\boldsymbol{x},t)\ - tv^i(\boldsymbol{x},t)$, $\delta_{ij} v^i(\boldsymbol{x},t)\ x^j$, respectively.

Corresponding to the complete generator (3.1) , we obtain:

$$\begin{cases} \bar{\delta}_0\ t &= 0 \\ \bar{\delta}_0 x^i &= \varepsilon(t)\dfrac{\delta^{ij} p_j}{m} + \eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t) \quad , \end{cases} \tag{3.4}$$

and

$$\begin{cases} \bar{\delta}_0 p_i &= p_k \dfrac{\partial}{\partial x^i}[\eta^k(\boldsymbol{x},t) - t\ v^k(\boldsymbol{x},t)] + m\dfrac{\partial}{\partial x^i}[\delta_{lk} x^l v^k(\boldsymbol{x},t)] \\ \bar{\delta}_0 \bar{H} &= \dfrac{1}{m}\delta^{ij} p_i \bar{\delta} p_j \\ \bar{\delta}_0 \bar{J}_i &= c_{ij}{}^k p_k [\eta^j(\boldsymbol{x},t) - t\ v^j(\boldsymbol{x},t)] - c_{ij}{}^k x^j p_r \dfrac{\partial}{\partial x^k}[\eta^r(\boldsymbol{x},t) - t\ v^r(\boldsymbol{x},t)] \\ & \quad - m c_{ij}{}^k x^j \dfrac{\partial}{\partial x^k}[\delta_{lm} x^l v^m(\boldsymbol{x},t)] \\ \bar{\delta}_0 \bar{K}_i &= \varepsilon(t) p_i + t p_k \dfrac{\partial}{\partial x^i}[\eta^k(\boldsymbol{x},t) - t\ v^k(\boldsymbol{x},t)] \\ & \quad + m\delta_{ij}[\eta^j(\boldsymbol{x},t) - t\ v^j(\boldsymbol{x},t)] - m\ t\dfrac{\partial}{\partial x^i}[\delta_{lm} x^l v^m(\boldsymbol{x},t)] \quad . \end{cases} \tag{3.5}$$

Finally, using $p_i = m\delta_{ij}\dot{x}^j$, we obtain

$$\left\{\begin{array}{lcl} \delta_0\, t & = & 0 \\ \delta_0 x^i & = & \varepsilon(t)\dot{x}^i + \eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t) \\ \delta_0 \dot{x}^i & = & \dfrac{d}{dt}[\varepsilon(t)\dot{x}^i] + \dot{x}^k \dfrac{\partial}{\partial x^k}[\eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t)] \\ & & +\dfrac{\partial}{\partial t}[\eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t)] \quad . \end{array}\right. \tag{3.6}$$

Eqs.(3.6) reduce to Eqs.(2.4) in the limit of global flat symmetry and they can be taken as a *definition* of the *Galilei gauge transformations* in configuration space. Notice that the *quasi-invariance* of the Lagrangian under the global flat transformations is now broken and that $\bar{\delta}_0 p_i|_{p_i = m\delta_{ij}\dot{x}^j}$ is not equal to $\delta_0[m\delta_{ij}\dot{x}^j]$.

On the other hand, within the re-parameterization invariant picture, corresponding to the complete generator (3.2), we get:

$$\left\{\begin{array}{lcl} \hat{\bar{\delta}}_0\ \lambda & = & 0 \\ \hat{\bar{\delta}}_0\ t(\lambda) & = & -\varepsilon(t(\lambda)) \\ \hat{\bar{\delta}}_0 x^i(\lambda) & = & \eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t) \\ \hat{\bar{\delta}}_0 t'(\lambda) & = & -t'\ \dfrac{d\varepsilon(t(\lambda))}{dt} \\ \hat{\bar{\delta}}_0 x'^i(\lambda) & = & x'^k \dfrac{\partial}{\partial x^k}[\eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t)] + t' \dfrac{\partial}{\partial t}[\eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t)]\ , \end{array}\right. \tag{3.7}$$

and

$$\left\{\begin{array}{lcl} \hat{\bar{\delta}}_0 \hat{p}_i & = & -\hat{p}_k \dfrac{\partial}{\partial x^i}[\eta^k(\boldsymbol{x},t) - t\ v^k(\boldsymbol{x},t)] - m\dfrac{\partial}{\partial x^i}[\delta_{lk} x^l v^k(\boldsymbol{x},t)] \\ \hat{\bar{\delta}}_0 \hat{E} & = & \hat{E}\dfrac{d\varepsilon(t)}{dt} + \hat{p}_i \dfrac{\partial}{\partial t}[\eta^i(\boldsymbol{x},t) - t\ v^i(\boldsymbol{x},t)] + m\dfrac{\partial}{\partial t}[\delta_{ij} x^i v^j(\boldsymbol{x},t)] \\ \hat{\bar{\delta}}_0 \hat{\mathcal{J}}_i & = & c_{ij}{}^k \hat{p}_k[\eta^j(\boldsymbol{x},t) - t\ v^j(\boldsymbol{x},t)] - c_{ij}{}^k x^j \hat{p}_r \dfrac{\partial}{\partial x^k}[\eta^r(\boldsymbol{x},t) - t\ v^r(\boldsymbol{x},t)] \\ & & -mc_{ij}{}^k x^j \dfrac{\partial}{\partial x^k}[\delta_{lm} x^l v^m(\boldsymbol{x},t)] \\ \hat{\bar{\delta}}_0 \hat{\mathcal{K}}_i & = & \varepsilon(t)\hat{p}_i + t\hat{p}_k \dfrac{\partial}{\partial x^i}[\eta^k(\boldsymbol{x},t) - t\ v^k(\boldsymbol{x},t)] \\ & & +m\delta_{ij}[\eta^j(\boldsymbol{x},t) - t\ v^j(\boldsymbol{x},t)] - mt\dfrac{\partial}{\partial x^i}[\delta_{lm} x^l v^m(\boldsymbol{x},t)] \quad . \end{array}\right. \tag{3.8}$$

In conclusion, from Eq.(3.6), we see that $\boldsymbol{x}(t)$ undergoes a time-dependent general Euclidean coordinate transformation (although deformed by an effect from Galilei boosts) plus a velocity-dependent transformation induced by the condition $\bar{\delta}_0 t = 0$. On the other hand, from Eq.(3.7), we see that $t(\lambda)$ undergoes a transformation which does not involve the Euclidean coordinates and, simultaneously, $\boldsymbol{x}(\lambda)$ undergoes a time-dependent general Euclidean coordinate transformation (still deformed by an effect from Galilei boosts). Being a consequence of the absolute nature of the Newtonian time, these features do not appear in the *gauging* of the Poincaré realization corresponding to the free relativistic mass-point with Lagrangian $L_M = -m\sqrt{\eta_{\mu\nu} x'^{\mu}(\lambda) x'^{\nu}(\lambda)}$ (see, for instance, [14]). Actually, the *gauging* of the Poincaré transformations induces a general coordinate transformation of $x^{\mu}(\lambda)$ which is indistinguishable from the simple *gauging* of the space-time translations alone. As well-known, the *gauging* of the Lorentz transformations is made evident by means of the "soldering procedure" which amounts to say that the vectors belonging to the tangent bundle of the curved space-time must transform as four vectors under "local" Lorentz transformation. The "soldering" is done by means of a set of *vierbeins* $\mathbf{E}^{A}_{\mu}(x)$ so that the flat transformation $x'^{\mu} \to x'^{\nu} + \varepsilon^{\mu}_{\cdot\nu} x'^{\mu}$ (where $\varepsilon^{\mu}_{\cdot\nu}$ are the parameters of the Lorentz transformations) is replaced by $\mathbf{E}^{A}_{\mu}(x) x'^{\mu} \to \mathbf{E}^{A}_{\mu}(x) x'^{\mu} + \varepsilon^{\mathcal{S}}_{\cdot B} \mathbf{E}^{A}_{\mu}(x) x'^{\mu}$.

We want to discuss now the problem of the invariance (possibly *quasi-invariance*) of the Lagrangian with respect to the local Galilei transformations just defined. In the Newtonian case, unlike the relativistic one, the absolute nature of time prevents us from using the standard "soldering" procedure, if not for the Euclidean subalgebra generated by $\hat{\mathcal{P}}_k$ and $\hat{\mathcal{J}}_i$. Actually, while the effect of the space rotations in the term $\eta^i(\boldsymbol{x}, t)$ is indistinguishable from the effect generated in it by space translations, the effect of "local" space rotations and translations can be distinguished for three-vectors by introducing the "soldering" with *dreibeins* $\mathbf{E}^a_i(\boldsymbol{x}, t)$. Notice, however, that further complications arise here from the fact that the general Euclidean transformations are time-dependent and that time translations and Galilei boosts introduce new terms into the transformation of the velocity. Accordingly, the rule for the transition from the global flat transformation of three-velocity to its general transformation is obtained by defining $\omega^a(\boldsymbol{x}, t) = \mathbf{E}^a_i(\boldsymbol{x}, t)\omega^i(\boldsymbol{x}, t)$, and by imposing, within the two alternative pictures introduced above, the following transformation properties

$$\left\{ \begin{array}{rcl} \delta_0\ \dot{x}^i & = & \dfrac{d}{dt}[\varepsilon \dot{x}^i] + c_{ij}{}^{k} \omega^j \dot{x}^k - v^i \\ \Rightarrow\ \ \delta_{0[t]}[\mathbf{E}^a_i \dot{x}^i] & = & \dfrac{d}{dt}[\varepsilon \mathbf{E}^a_i \dot{x}^i] + c_{bc}{}^{a} \omega^b \mathbf{E}^c_i \dot{x}^i + [?] \end{array} \right. \tag{3.9}$$

$$\left\{\begin{array}{rcl} \hat{\delta}_0\; x'^i & = & c_{jk}{}^i\omega^j x'^k - v^i t' \\ \Rightarrow\; \hat{\delta}_{0[\lambda]}[\mathbf{E}^a_i x'^i] & = & c_{bc}{}^a\omega^b\mathbf{E}^c_i x'^i + [?]\;\;, \end{array}\right. \tag{3.10}$$

where the question marks in Eqs.(3.9) and (3.10) stand for possible terms connected to time translation and Galilei boosts: this point will be settled later on.

By means of the invertible *dreibeins* $\mathbf{E}^a_i$, a Euclidean metric in the three-space is naturally introduced in the form

$$\delta_{ij} \Rightarrow g_{ij} \equiv \delta_{ab}\mathbf{E}^a_i\mathbf{E}^b_j \;\;, \tag{3.11}$$

while the inverse metric can be likewise defined in term of the inverse *dreibeins* $\mathbf{H}^i_a$ ($\mathbf{H}^i_a\mathbf{E}^b_i = \delta^a_b$, $\mathbf{H}^i_a\mathbf{E}^a_j = \delta^i_j$) as:

$$\delta^{ij} \rightarrow g^{ij} \equiv \delta^{ab}\mathbf{H}^i_a\mathbf{H}^j_b \;\;,\;\; g^{ij}g_{jk} = \delta^i_k \;\;. \tag{3.12}$$

Let us now discuss, within the standard picture, the issue of the invariance of the Lagrangian under the newly introduced local transformations, by considering first the local time-independent Euclidean transformations $\delta^{[e]}_0$ ($\varepsilon = 0$, $v^i = 0$, $\omega^i = \omega^i(\boldsymbol{x})$, $\varepsilon^i = \varepsilon^i(\boldsymbol{x})$); then pure time transformations $\delta^{[t]}_0$ ($\varepsilon = \varepsilon(t)$, $v^i = 0$, $\omega^i = 0$, $\varepsilon^i = 0$), and the combination of both $\delta^{[e]}_0$ and $\delta^{[t]}_0$, say $\delta^{[et]}_0$ ($\varepsilon = \varepsilon(t)$, $v^i = 0$, $\omega^i = \omega^i(\boldsymbol{x})$, $\varepsilon^i = \varepsilon^i(\boldsymbol{x})$); finally the most general transformations including local time-dependent Euclidean transformations and Galilei boosts.

i) In the case of local time-independent Euclidean transformations $\delta^{[e]}_0$, defined by $\varepsilon = 0$, $v^i = 0$, $\omega^i = \omega^i(\boldsymbol{x})$, $\varepsilon^i = \varepsilon^i(\boldsymbol{x})$, we have:

$$\begin{array}{rcl} \delta^{[e]}_0\dot{x}^i & = & \dot{x}^k\dfrac{\partial}{\partial x^k}[\varepsilon^i(\boldsymbol{x}) + c_{jl}{}^i\omega^j(\boldsymbol{x})x^l] \\ & = & \dot{x}^k\dfrac{\partial\eta^i_{[e]}(\boldsymbol{x})}{\partial x^k} \;\;, \end{array} \tag{3.13}$$

where $\eta^i_{[e]}(\boldsymbol{x}) = \varepsilon^i(\boldsymbol{x}) + c_{jl}{}^i\omega^j(\boldsymbol{x})x^l$ is the restriction of $\eta^i(\boldsymbol{x},t)$ to time independent Euclidean transformation. Then, by imposing

$$\delta^{[e]}_{0[t]}[\mathbf{E}^a_i\dot{x}^i] = c_{bc}{}^a\omega^b(\boldsymbol{x})[\mathbf{E}^c_i\dot{x}^i]\;, \tag{3.14}$$

it follows:

$$\begin{array}{rcl} \delta^{[e]}_{0[t]}\mathbf{E}^a_i(\boldsymbol{x},t) & = & \delta^{[e]}_0\mathbf{E}^a_i(\boldsymbol{x},t) + \dfrac{\partial\mathbf{E}^a_i(\boldsymbol{x},t)}{\partial x^k}\delta^{[e]}_0 x^k \\ & = & -\dfrac{\partial\eta^i_{[e]}(\boldsymbol{x})}{\partial x^k}\mathbf{E}^a_i + c_{bc}{}^a\omega^b(\boldsymbol{x})\mathbf{E}^c_i \;\;, \end{array} \tag{3.15}$$

i.e.

$$\delta^{[e]}_{0[t]}\mathbf{E}^a_i = -\frac{\partial \eta^i_{[e]}(\boldsymbol{x})}{\partial x^k}\mathbf{E}^a_i + c_{bc}{}^a\omega^b(\boldsymbol{x})\mathbf{E}^c_i - \mathbf{E}^a_{i,k}\eta^k_{[e]}(\boldsymbol{x}) \ . \tag{3.16}$$

in agreement to the conventions established in Eqs.(2.13). It is then immediately seen that the invariance of the Lagrangian under time-independent local Euclidean transformation can be recovered by means of the substitution:

$$\dot{x}^i \Rightarrow \mathbf{E}^a_i\dot{x}^i \quad , \tag{3.17}$$

which leads to

$$\begin{aligned} L_M = \tfrac{1}{2}m\delta_{ij}\dot{x}^i\dot{x}^j \quad \Rightarrow L^{g_{[e]}}_M &\equiv \tfrac{1}{2}m\delta_{ab}\mathbf{E}^a_i\dot{x}^i\mathbf{E}^b_j\dot{x}^j \\ &= \tfrac{1}{2}mg_{ij}\dot{x}^i\dot{x}^j \ . \end{aligned} \tag{3.18}$$

In fact, since the transformation rules for the three-dimensional metric field (3.11) are given by:

$$\delta^{[e]}_{0[t]}g_{ij} = \delta^{[e]}_0 g_{ij} + g_{ij,k}\eta^k_{[e]}(\boldsymbol{x}) = -\frac{\partial \eta^k_{[e]}(\boldsymbol{x})}{\partial x^i}g_{kj} - \frac{\partial \eta^k_{[e]}(\boldsymbol{x})}{\partial x^j}g_{ik} \quad , \tag{3.19}$$

(which are the correct infinitesimal transformation properties of a covariant second-rank three-dimensional tensor), it follows:

$$\delta^{[e]}_{0[t]}L^{g_{[e]}}_M = 0 \quad . \tag{3.20}$$

Therefore, the preliminary guess for the *gauging* of the Galilei transformation (see Eqs. 3.1, 3.4-3.6, or 3.2, 3.7 and 3.8) is made geometrically consistent by replacing the flat metric δ_{ij} and δ^{ij} in those formulae by g_{ij} and g^{ij}, respectively.

Within the re-parameterization invariant picture, the invariance is recovered by the corresponding substitution,

$$x'^i \Rightarrow \mathbf{E}^a_i x'^i \tag{3.21}$$

which leads to

$$\hat{L}_M(\lambda) \Rightarrow \hat{L}^{g_{[e]}}_M(\lambda) = \frac{1}{2}m\frac{g_{ij}x'^i(\lambda)x'^j(\lambda)}{t'(\lambda)} \ . \tag{3.22}$$

In fact, from Eq.(3.19) with $\delta^{[e]}_{0[t]}$ formally replaced by $\hat{\delta}^{[e]}_{0[\lambda]}$, it follows:

$$\hat{\delta}^{[e]}_{0[\lambda]}\hat{L}^{[e]}_M = 0 \ . \tag{3.23}$$

ii) Consider now the pure time-translations $\hat{\delta}_0^{[t]}$. For the sake of simplicity, we shall confine to the re-parameterization invariant picture. We have:

$$\left\{ \begin{array}{lcl} \hat{\delta}_0^{[t]}\, t & = & -\epsilon(t) \\ \hat{\delta}_0^{[t]} x^i & = & 0 \quad . \end{array} \right. \tag{3.24}$$

It is immediate to see that, by means of a *einbein* substitution of the form

$$t' \Rightarrow \Theta(t)t' \quad , \tag{3.25}$$

the Lagrangian:

$$\hat{L}_M^{g[t]}(\lambda) \equiv \frac{1}{2}m\frac{\delta_{ij}x'^i(\lambda)x'^j(\lambda)}{\Theta(t)t'(\lambda)} \quad , \tag{3.26}$$

is invariant under the transformations $\hat{\delta}_0^{[t]}$, provided we impose:

$$\hat{\delta}_{0[\lambda]}^{[t]}[\Theta(t)t'] = 0 \quad , \tag{3.27}$$

which, in turn, gives:

$$\hat{\delta}_{0[\lambda]}^{[t]}\Theta(t) = \hat{\delta}_0^{[t]}\Theta(t) - \varepsilon(t)\frac{d\Theta(t)}{dt} = \dot{\varepsilon}(t)\Theta(t) \quad . \tag{3.28}$$

iii) We can summarize the results found up to now in the re-parameterization invariant picture, by saying that the modified matter Lagrangian

$$\hat{L}_M^{g[et]}(\lambda) \equiv \frac{1}{2}m\frac{g_{ij}x'^i(\lambda)x'^j(\lambda)}{\Theta(t)t'(\lambda)} \quad , \tag{3.29}$$

is strictly invariant under the local transformations:

$$\left\{ \begin{array}{lclcl} \hat{\delta}_0^{[et]} t & = & \hat{\delta}_0^{[t]} t & = & -\epsilon(t) \\ \hat{\delta}_0^{[et]} x^i & = & \hat{\delta}_0^{[e]} x^i & = & \eta^i_{[e]}(\boldsymbol{x}) \\ \hat{\delta}_0^{[et]} t' & = & \hat{\delta}_0^{[t]} t' & = & -\dot{\varepsilon}(t)t' \\ \hat{\delta}_0^{[et]} x'^i & = & \hat{\delta}_0^{[e]} x'^i & = & x'^k\dfrac{\partial\eta^i_{[e]}(\boldsymbol{x})}{\partial x^k} \quad , \end{array} \right. \tag{3.30}$$

if we adopt the following transformation rules for the fields:

$$\left\{ \begin{array}{lclcl} \hat{\delta}_{0[\lambda]}^{[et]}\Theta(t) & = & \hat{\delta}_{0[\lambda]}^{[t]}\Theta(t) & = & \dot{\varepsilon}(t)\Theta(t) \\ \hat{\delta}_{0[\lambda]}^{[et]}\mathbf{E}_i^a & = & \hat{\delta}_{0[\lambda]}^{[e]}\mathbf{E}_i^a & = & -\dfrac{\partial\eta^i_{[e]}(\boldsymbol{x})}{\partial x^k}\mathbf{E}_i^a + c_{bc}{}^a\omega^b(\boldsymbol{x})\mathbf{E}_i^c \\ \hat{\delta}_{0[\lambda]}^{[et]} g_{ij} & = & \hat{\delta}_{0[\lambda]}^{[e]} g_{ij} & = & -\dfrac{\partial\eta^k_{[e]}(\boldsymbol{x})}{\partial x^i}g_{kj} - \dfrac{\partial\eta^k_{[e]}(\boldsymbol{x})}{\partial x^j}g_{ik} \quad . \end{array} \right. \tag{3.31}$$

iv) Finally, we have to show that the invariance (possibly *quasi-invariance*) of the matter Lagrangian under the most general transformations, including local time-dependent Euclidean transformations and Galilei boosts, can be recovered by introducing *four* additional external fields, say $A_0(\boldsymbol{x},t)$, $A_i(\boldsymbol{x},t)$. Within the re-parameterization invariant picture, our scope will be achieved if we succeed in defining a new matter Lagrangian $\hat{L}^g_M$ such that:
a) under the *gauge* transformations (3.7) and the corresponding transformations induced on all the additional fields, be *quasi-invariant* in the form:

$$\hat{\delta}_{0[\lambda]}\hat{L}^g_M = \frac{d\mathcal{F}}{d\lambda} \quad , \tag{3.32}$$

b) its *global flat* limit, together with that of its transformation properties, coincide with the expression (2.23) and (2.32), respectively.

First of all, in order to reproduce the *cocycle* term of Eq.(2.32) in the global flat limit, $\mathcal{F}$ will be chosen as $\mathcal{F} = -mg_{ij}v^i x^j$. Then, the matter Lagrangian will be defined as

$$\hat{L}^g_M(\lambda) = \frac{1}{\Theta t'}\frac{m}{2}\left[g_{ij}x'^i x'^j + 2A_i x'^i t' + 2A_0 t' t'\right] \quad . \tag{3.33}$$

Second, since we have to preserve the transformation properties of $\mathbf{E}^a_i$, g_{ij} and Θ already established for the case $v^i(\boldsymbol{x},t) = 0$ in Eqs.(3.31), we shall assume

$$\left\{\begin{aligned} \hat{\delta}_{0[\lambda]}\Theta &= \hat{\delta}_0\Theta - \epsilon(t)\frac{d\Theta(t)}{dt} \\ &= \dot{\epsilon}(t)\Theta(t) \\ \hat{\delta}_{0[\lambda]}\mathbf{E}^a_i &= \hat{\delta}_0\mathbf{E}^a_i + \mathbf{E}^a_{i,k}\tilde{\eta}^k(\boldsymbol{x},t) - \frac{\partial\mathbf{E}^a_i}{\partial t}\varepsilon(t) \\ &= c_{bc}{}^a\omega^b(\boldsymbol{x},t)\mathbf{E}^c_i - \frac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}\mathbf{E}^a_k \end{aligned}\right. \tag{3.34}$$

$$\Rightarrow \quad \begin{aligned} \hat{\delta}_{0[\lambda]}g_{ij} &= \hat{\delta}_0 g_{ij} + g_{ij,k}\tilde{\eta}^k(\boldsymbol{x},t) - \frac{\partial g_{ij}}{\partial t}\varepsilon(t) \\ &= -\frac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}g_{kj} - \frac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}g_{kj} \ , \end{aligned}$$

where we have introduced the notation $\tilde{\eta}^k(\boldsymbol{x},t) \equiv \eta^k(\boldsymbol{x},t) - tv^k(\boldsymbol{x},t)$. Then, the *quasi-invariance* (3.32) of the matter Lagrangian (3.33) is guaranteed if the additional fields

$A_0(\boldsymbol{x},t)$, $A_i(\boldsymbol{x},t)$ transform as follows:

$$\begin{cases} \hat{\delta}_{0[\lambda]}A_0 &= \hat{\delta}_0 A_0 - \varepsilon\dfrac{\partial A_0}{\partial t} + A_{i,j}\tilde{\eta}^j \\ &= 2\dot{\varepsilon}A_0 - A_i\dfrac{\partial\tilde{\eta}^i}{\partial t} + \Theta\dfrac{\partial\mathcal{F}}{\partial t} \\ \hat{\delta}_{0[\lambda]}A_i &= \hat{\delta}_0 A_i - \varepsilon\dfrac{\partial A_i}{\partial t} + A_{i,j}\tilde{\eta}^j \\ &= \dot{\varepsilon}A_i - A_j\dfrac{\partial\tilde{\eta}^j}{\partial x^i} - g_{ij}\dfrac{\partial\tilde{\eta}^j}{\partial t} + \Theta\dfrac{\partial\mathcal{F}}{\partial x^i} \quad . \end{cases} \tag{3.35}$$

Notice that the fields $\mathbf{E}^a_i$, g_{ij}, Θ, A_0 and A_i have the global flat Galilean limits δ^a_i, δ_{ij}, 1, 0 and 0, respectively. Here, the A_i's play the role of the components $\mathbf{E}^0_i(x)$ of the relativistic *vierbeins*, while the two fields Θ and A_0 correspond to a splitting of the *vierbein* component $\mathbf{E}^0_0(x)$, as it will be clear later on.

In a similar way, it can be seen that, within the standard picture, the Lagrangian

$$L^g_M(t) = \frac{1}{\Theta}\frac{m}{2}\left[g_{ij}\dot{x}^i\dot{x}^j + 2A_i\dot{x}^i + 2A_0\right] \quad , \tag{3.36}$$

is *quasi-invariant* under the field transformation rules (3.34), (3.35), with $\delta_0 = \hat{\delta}_0$, and the mass-point coordinate transformations given by (3.6), in the sense that we have

$$\delta_{0[t]}L^g_M = \frac{d\mathcal{F}}{dt} + \varepsilon\frac{d[L^g_M]}{dt} \ , \tag{3.37}$$

so that

$$\Delta\mathcal{S}^g_M = \int_{t_1}^{t_2} dt\left[\frac{d\mathcal{F}}{dt} + \frac{d[\varepsilon L^g_M]}{dt}\right] = 0 \quad . \tag{3.38}$$

The condition for the invariance of the theory under the *gauged* Galilei transformations can now be easily formulated also in the Hamiltonian formalism. From Eq.(3.33), we get the following expressions for energy and linear momentum

$$\begin{cases} \hat{E} &= \dfrac{\partial\hat{L}^g_M}{\partial t'} = \dfrac{1}{\Theta t'^2}\dfrac{m}{2}g_{ij}x'^i x'^j - \dfrac{m}{\Theta}A_0 \\ \hat{p}_i &= \dfrac{\partial\hat{L}^g_M}{\partial x'^i} = \dfrac{m}{\Theta t'}\left(g_{ij}x'^j + A_i t'\right) \quad , \end{cases} \tag{3.39}$$

while the first-class constraint (2.27) becomes

$$\hat{\chi}_g = \frac{1}{\Theta}\left[\hat{E} + \frac{m}{\Theta}A_0\right] - \frac{g^{ij}}{2m}\left[\hat{p}_i - \frac{m}{\Theta}A_i\right]\left[\hat{p}_j - \frac{m}{\Theta}A_j\right] \quad . \tag{3.40}$$

Since, under the transformation (3.8) (with $\delta_{ij} \Rightarrow g_{ij}$), we have:

$$\begin{aligned}
\hat{\bar{\bar{\delta}}}_{0[\lambda]}\hat{\chi} \;&= \frac{1}{\Theta}\hat{\bar{\bar{\delta}}}_0\hat{E} - \frac{1}{\Theta^2}\left[\hat{E} + \frac{m}{\Theta}A_0\right]\hat{\delta}_{0[\lambda]}\Theta + \frac{1}{\Theta^2}m\hat{\delta}_{0[\lambda]}A_0 \\
&- \frac{1}{2m}\hat{\delta}_{0[\lambda]}g^{ij}\left[\hat{p}_i - \frac{m}{\Theta}A_i\right]\left[\hat{p}_j - \frac{m}{\Theta}A_j\right] \\
&- \frac{g^{ij}}{m}\left[\hat{p}_i - \frac{m}{\Theta}A_i\right]\left[\hat{\bar{\bar{\delta}}}_0\hat{p}_j - \frac{m}{\Theta}\hat{\delta}_{0[\lambda]}A_j + \frac{m}{\Theta^2}A_j\hat{\delta}_{0[\lambda]}\Theta\right] \quad ,
\end{aligned} \tag{3.41}$$

the invariance of the constraint

$$\hat{\bar{\bar{\delta}}}_{0[\lambda]}\hat{\chi}_g = 0 \quad , \tag{3.42}$$

which is the Hamiltonian analogue to the invariance of the Lagrangian, is guaranteed if the fields transform according to

$$\begin{cases}
\hat{\delta}_{0[\lambda]}\ \Theta &= \dot{\epsilon}(t)\Theta(t) \\
\hat{\delta}_{0[\lambda]}g_{ij} &= -\dfrac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}g_{kj} - \dfrac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}g_{kj} \\
\hat{\delta}_{0[\lambda]}A_0 &= 2\dot{\varepsilon}A_0 - A_i\dfrac{\partial\tilde{\eta}^i}{\partial t} - \Theta\dfrac{\partial}{\partial t}\left[g_{ij}v^i x^j\right] \\
\hat{\delta}_{0[\lambda]}A_i &= \dot{\varepsilon}A_i - A_j\dfrac{\partial\tilde{\eta}^j}{\partial x^i} - g_{ij}\dfrac{\partial\tilde{\eta}^j}{\partial t} - \Theta\dfrac{\partial}{\partial x^i}\left[g_{ij}v^i x^j\right] \quad .
\end{cases} \tag{3.43}$$

Note that since

$$\delta f(\boldsymbol{x}(t),t) = \hat{\delta}_{0[\lambda]}f(\boldsymbol{x}(t(\lambda)),t(\lambda)) \quad , \tag{3.44}$$

Eqs.(3.43) can be easily adapted to the standard picture.

Let us remark in addition that were it not for the presence of the *einbein* Θ, the modified Hamiltonian constraints (3.40) could have been made invariant only in the *week* sense $\hat{\bar{\bar{\delta}}}_{0[\lambda]}\hat{\chi}_g = \varepsilon\hat{\chi}_g \simeq 0$.

Finally, the *cocycle* term which appeared in a generic form in the transformations (3.35) of the fields A_0 and A_i, is now *explicitly determined* (up to a constant) in a form which reproduces the standard expression in the global flat Galilean limit. This is due to the fact that the Hamiltonian formalism of the reparemetrization invariant scheme requires a first class constraint which, in absence of external fields, says that the Galilei Casimir invariant representing the internal energy ($E - \frac{1}{2m}\delta^{ij}p_i p_j$) is equal to zero.

In presence of external fields, the constraints given by Eqs.(3.40), says again that the Casimir invariant vanishes. This is clearly a consequence of the fact that time is absolute and of the fact that preservation of the constraints (i.e. of the vanishing of the Casimir invariant) is possible only provided the projective realization is taken into account through the *cocycle* terms in Eq.(3.43). The analogue of this phenomenon in the standard Lagrangian picture is expressed by the fact that the identification of the total energy in different coordinate systems connected by general Galilean coordinate transformation requires explicitly the appearance of the *cocycle* term within the transformation rules of the relevant quantities. On the other hand it is obvious that the *cocycle* term is not associated to actual forces since it appears in the variation of the Lagrangian as a total derivative irrelevant for the equations of motion.

4 Dynamics of a mass-point in the external "gauge" fields

We want to discuss now the equations of motion of the mass-point acted upon, as a *test particle*, by the newly introduced compensating external fields. For the sake of simplicity, we will use the Lagrangian in the standard picture:

$$L^g_M(t) = \frac{m}{\Theta}\left[\frac{1}{2} g_{ij}\dot{x}^i\dot{x}^j + A_i\dot{x}^i + A_0\right] \quad . \tag{4.1}$$

Upon variation, we get the Euler-Lagrange equations:

$$\begin{aligned} \ddot{x}^i + \Gamma^i_{kl}\dot{x}^k\dot{x}^l &= -\frac{\dot{\Theta}}{\Theta}\left[\dot{x}^i + g^{ij}A_j\right] - g^{ij}\frac{\partial g_{jl}}{\partial t}\dot{x}^l \\ &\quad + g^{ij}\left[\frac{\partial A_0}{\partial x^j} - \frac{\partial A_j}{\partial t}\right] + g^{ij}\left[\frac{\partial A_l}{\partial x^j} - \frac{\partial A_j}{\partial x^l}\right]\dot{x}^l \quad , \end{aligned} \tag{4.2}$$

where:

$$\Gamma^l_{ij} = \left\{ \begin{matrix} l \\ i\ j \end{matrix} \right\} = \frac{1}{2}\mathbf{H}^l_a[\mathbf{E}^a_{i,j} + \mathbf{E}^a_{j,i}] \quad , \tag{4.3}$$

is the three-dimensional metric affinity of g_{ij}.

Let us stress that the function $\Theta(t)$ which appears in the Lagrangian (4.1) and in the equations of motion (4.2) has no real dynamical content. In fact, by redefining the evolution parameter t and the field A_0, A_i according to

$$\begin{cases} T(t) & \equiv \int_0^t d\tau \Theta(\tau) \\ \tilde{A}_0 & \equiv \dfrac{A_0}{\Theta^2} \\ \tilde{A}_i & \equiv \dfrac{A_i}{\Theta} \quad , \end{cases} \tag{4.4}$$

it follows:

$$\begin{aligned} \frac{d^2x^i}{dT^2} + \Gamma^i_{kl}\frac{dx^k}{dT}\frac{dx^l}{dT} &= -g^{ij}\frac{\partial g_{jl}}{\partial T}\frac{dx^l}{dT} + g^{ij}\left[\frac{\partial \tilde{A}_l}{\partial x^j} - \frac{\partial \tilde{A}_i}{\partial x^l}\right]\frac{dx^l}{dT} \\ &\quad + g^{ij}\left[\frac{\partial \tilde{A}_0}{\partial x^j} - \frac{\partial \tilde{A}_i}{\partial T}\right] \quad . \end{aligned} \tag{4.5}$$

In order to gain a better physical insight, we will discuss the standard Newtonian gravitational problem in general Galilean coordinates. Before doing that, let us reproduce in our notations the Kuchař's classification of the relevant reference frames obtained by passive coordinate transformations:

1) *non rotating observers*: $\Theta = 1$, $A_i = 0$;

2) *rigid observers*, $\Theta = 1$, $g_{ij} = \delta_{ij}$ (an example belonging to this class is provided by eqs.(4.19) below; Wheeler' definition of *Galilean observer* is a subcase of *rigid observers*, see eq.(4.22);

3) *freely-falling observers*: $\Theta = 1$, $A_0 = 0$

4) *Gaussian* (freely falling, non rotating) *observer*: $\Theta = 1$, $A_i = 0$, $A_0 = 0$ (these are the analogues of the *Fermi-Walker observers* in general relativity);

5) *Galilean observers* (rigid, non rotating): $\Theta = 1$, $g_{ij} = \delta_{ij}$, $A_i = 0$, $A_0 = -\varphi$: they are called *absolute Galilean observers* by Wheeler [15] and *inertial* by Levi-Civita [16];

6) *inertial observers* (rigid, freely falling, non rotating): $\Theta = 1$, $g_{ij} = \delta_{ij}$, $A_i = 0$, $A_0 = 0$;

Let $[y^a, t]$ be the coordinates used by a *Galilean observer* (rigid and non-rotating). Newton's equations can be written:

$$m\frac{d^2y^a}{dt^2} = -m\delta^{ab}\frac{\partial \varphi}{\partial y^b} \ , \tag{4.6}$$

where $\varphi(\boldsymbol{y}, t)$ is the gravitational potential satisfying the Poisson equation ($\Delta = \delta^{ij}\partial_i\partial_j$)

$$\Delta\varphi(\boldsymbol{z}, t) = 4\pi G m \delta[\boldsymbol{z} - \boldsymbol{y}(t)] \ \ . \tag{4.7}$$

This situation corresponds to:

$$\Theta = 1, \qquad g_{ij} = \delta_{ij}, \qquad A_0 = -\varphi, \qquad A_i = 0\ . \tag{4.8}$$

Eqs.(4.6) and (4.7) are form-invariant under the group of Galilei transformations (2.3) that connect *Galilean observers*. In particular, the gravitational potential is a *scalar* under these transformations (see for example Wheeler [15] §12.17).

Let us perform now a passive transformation to an arbitrary coordinate system (T, ξ^a):

$$\left\{ \begin{array}{ll} t & \rightarrow T(t) \\ y^a & \rightarrow \xi^a(\boldsymbol{x}(t), t) \quad . \end{array} \right. \tag{4.9}$$

We have:

$$\begin{aligned} \frac{d^2\xi^a}{dT^2} \ = \ & \frac{1}{\dot{T}^2}\left[\frac{\partial \xi^a}{\partial x^i}\ddot{x}^i\right] + \frac{\ddot{T}}{\dot{T}^t}\left[\frac{\partial \xi^a}{\partial x^i}\dot{x}^i + \frac{\partial \xi^a}{\partial t}\right] \\ & + \frac{1}{\dot{T}^2}\left[\frac{\partial^2 \xi^a}{\partial x^i \partial x^j}\dot{x}^i\dot{x}^j + 2\frac{\partial^2 \xi^a}{\partial x^i \partial t}\dot{x}^i + \frac{\partial^2 \xi^a}{\partial t \partial t}\right] \quad , \end{aligned} \tag{4.10}$$

and, after some algebraic manipulations, using Eq.(4.3),

$$\begin{aligned} \ddot{x}^i + \frac{\partial x^i}{\partial y^a}\frac{\partial^2 \xi^a}{\partial x^k \partial x^l}\dot{x}^i\dot{x}^j \ = \ & -\frac{\partial x^i}{\partial y^a}\left[2\frac{\partial^2 \xi^a}{\partial x^i \partial t}\dot{x}^i + \frac{\partial^2 \xi^a}{\partial t \partial t}\right] \\ & -\frac{\partial x^i}{\partial y^a}\left[\frac{\ddot{T}}{\dot{T}}\left[\frac{\partial \xi^a}{\partial x^i}\dot{x}^i + \frac{\partial \xi^a}{\partial t}\right]\right] \\ & -\frac{\partial x^i}{\partial y^a}\dot{T}^2\delta^{ab}\frac{\partial \varphi}{\partial y^b} \quad , \end{aligned} \tag{4.11}$$

where $\dot{x}$ means derivation respect the evolution parameter t. The identification of the equations (4.11) and (4.2) expresses the non-relativistic *equivalence principle* which is implicit in the application of the *gauge* technique to a space-time symmetry group. We have:

$$\left\{ \begin{array}{ll} \Theta & = \dot{T} \\ g_{ij} & = \delta_{ab}\dfrac{\partial \xi^a}{\partial x^i}\dfrac{\partial \xi^b}{\partial x^j} \quad ; \ \mathbf{E}^a_i = \dfrac{\partial \xi^a}{\partial x^i} \\ g^{ij} & = \delta^{ab}\dfrac{\partial x^i}{\partial \xi^a}\dfrac{\partial x^j}{\partial \xi^b} \quad ; \ \mathbf{H}^i_a = \dfrac{\partial x^i}{\partial y^a} \\ A_0 & = -\varphi\dot{T}^2 + \dfrac{1}{2}\delta_{ab}\dfrac{\partial \xi^a}{\partial t}\dfrac{\partial \xi^b}{\partial t} \\ A_i & = \delta_{ab}\dfrac{\partial \xi^a}{\partial x^i}\dfrac{\partial \xi^b}{\partial t} \quad . \end{array} \right. \tag{4.12}$$

Note that

$$\tilde{A} = \frac{1}{\Theta^2}(A_0 - g^{ij}A_iA_j) = -\varphi \tag{4.13}$$

Then this last combination of the fields A_0 and A_i has to be identified with Newton's potential, while the A_i's are inertial fields.

Let us remark that, while the Poisson equation (4.7) can be rewritten in the new coordinates $z^b \rightarrow \xi^b(\boldsymbol{z}', t)$ in the form

$$\frac{1}{\sqrt{g}}\frac{\partial}{\partial z'^i}\left[\sqrt{g}g^{ij}\frac{\partial\varphi}{\partial z'^j}\right] = 4\pi Gm\delta[\boldsymbol{z}' - \boldsymbol{x}(t)] \quad , \tag{4.14}$$

this equation does not determine the gravitational potential since this latter is not a geometrical object and its functional form must depend on other fields. Therefore, the transformed Poisson equation (4.14) cannot be considered the equation for the field A_0 in an arbitrary reference frame and its integration cannot give rise to a term like $\frac{1}{2}\delta_{ab}\frac{\partial\xi^a}{\partial t}\frac{\partial\xi^b}{\partial t}$, as it should be according to Eq.(4.12). Instead, it is the quantity $\tilde{A}$ of Eqs.(4.13) that plays now the role of the Newton's potential φ. Then Eqs.(4.14) can be rewritten in the form

$$\frac{1}{\sqrt{g}}\frac{\partial}{\partial z'^i}\left[\sqrt{g}g^{ij}\frac{\partial\tilde{A}}{\partial z'^j}\right] = -4\pi Gm\delta[\boldsymbol{z}' - \boldsymbol{x}(t)] \quad . \tag{4.15}$$

The existence of the *cocycle* terms in the transformation rules of the potentials (see Eqs.(3.43)) does not invalidate the *non-relativistic equivalence principle* since, as shown at the end of the previous section, it affects only the definition of the total energy in different coordinate systems. The dynamics of a mass-point in the above external fields can be made Galilean-generally-covariant only if the second Galilean Casimir invariant $(E - P^2/2m)$ vanishes, as in the flat case.

Let us consider now various cases of passive coordinate transformations. For instance the kinematical group of passive coordinate transformations for the *rigid observers* (i.e., an arbitrary rigid but rotating frame). First, there are the *Galilean coordinate transformations* (according to Wheeler's terminology), which preserve the rigid and non-rotating character of the coordinates:

$$\left\{\begin{array}{lcllcl} t & \rightarrow & t - \varepsilon & , & \varepsilon & = & \text{cost.} \\ y^a & \rightarrow & \xi^a(\boldsymbol{x}, t) = \mathrm{R}^a_i x^i + \varepsilon^a(t) & , & \mathrm{R}^a_i & = & \text{cost.} \ . \end{array}\right. \tag{4.16}$$

We have now $\Theta = 1$, $g_{ij} = \delta_{ij}$, $A_0 = -\varphi + \frac{1}{2}\delta_{ab}\dot{\varepsilon}^a\dot{\varepsilon}^b$, $A_i = \delta_{ab}\mathrm{R}^a_i\dot{\varepsilon}^b$, and Eq.(4.6) is correspondingly modified.

These results can be obtained as a particular case ($\dot{\mathrm{R}}^a_i = 0$) of the passive coordinate transformation corresponding to an *arbitrary rigid* and *rotating* motion

$$\left\{ \begin{array}{ll} t & \to t - \varepsilon \\ y^a & \to \xi^a(\boldsymbol{x}, t) = \mathrm{R}^a_i(t)x^i + \varepsilon^a(t) \quad . \end{array} \right. \tag{4.17}$$

In this case we have:

$$\left\{ \begin{array}{ll} \dfrac{\partial \xi^a}{\partial x^j} & \equiv \mathbf{E}^a_j(\boldsymbol{x}, t) = \mathrm{R}^a_j(t) \\ \dfrac{\partial \xi^a}{\partial t} & = \dot{\mathrm{R}}^a_j(t)x^j + \dot{\varepsilon}^a(t) \\ \dfrac{\partial^2 \xi^a}{\partial x^j \partial t} & = \dot{\mathrm{R}}^a_j(t) \\ \dfrac{\partial^2 \xi^a}{\partial t^2} & = \ddot{\mathrm{R}}^a_j(t)x^j + \ddot{\varepsilon}^a(t) \quad , \end{array} \right. \tag{4.18}$$

and Eqs.(4.12) become:

$$\left\{ \begin{array}{lll} \Theta & = & 1 \\ g_{ij} & = & \delta_{ij} \; ; \; \dfrac{\partial g_{ij}}{\partial t} = 0 \; ; \; \Gamma^k_{ij} = 0 \\ A_0 & = & -\varphi + \dfrac{1}{2}\delta_{ab}[\dot{\mathrm{R}}^a_j x^j + \dot{\varepsilon}^a][\dot{\mathrm{R}}^b_j x^j + \dot{\varepsilon}^b] \\ A_i & = & \delta_{ab}\mathrm{R}^a_i[\dot{\mathrm{R}}^b_j x^j + \dot{\varepsilon}^b] \quad . \end{array} \right. \tag{4.19}$$

Then, the equations of motions take the form:

$$\begin{array}{rcl} \ddot{x}^l & = & \delta^{lk}\left\{ \dfrac{\partial A_k}{\partial t} - \dfrac{\partial A_0}{\partial x^k} + \left[\dfrac{\partial A_k}{\partial x^j} - \dfrac{\partial A_j}{\partial x^k} \right] \dot{x}^j \right\} \\ & = & \delta^{lk}\left\{ -\dfrac{\partial \phi}{\partial x^k} - \delta_{ab}\mathrm{R}^a_k\ddot{\varepsilon}^b - 2\delta_{ab}\mathrm{R}^a_k\dot{\mathrm{R}}^b_j\dot{x}^j \right. \\ & & \left. -\dfrac{1}{2}\delta_{ab}\left[\mathrm{R}^a_k\ddot{\mathrm{R}}^b_j + \ddot{\mathrm{R}}^a_k\mathrm{R}^b_j\right]x^j - \dfrac{1}{2}\delta_{ab}\left[\mathrm{R}^a_k\ddot{\mathrm{R}}^a_j - \ddot{\mathrm{R}}^a_k\mathrm{R}^a_j\right]x^j \right\} \quad . \end{array} \tag{4.20}$$

Since the angular velocity vector can be expressed as

$$\omega^k(t) = \frac{1}{2}\epsilon^{krs}\delta_{ab}\mathrm{R}^a_r(t)\dot{\mathrm{R}}^b_s(t) \quad , \tag{4.21}$$

the physical meaning of the various terms can be immediately identified as follows:

$$\begin{array}{rcll} m[\dfrac{\partial A_k}{\partial x^j} - \dfrac{\partial A_j}{\partial x^k}]\dot{x}^j & = & 2m\mathrm{R}^a_j(t)\dot{\mathrm{R}}^a_k(t)\dot{x}^j = 2m[\vec{\omega} \wedge \dot{\vec{x}}]_k & \text{(Coriolis force)} \\ m[\dfrac{\partial A_k}{\partial t} - \dfrac{\partial A_0}{\partial x^k}] & = & -m\dfrac{\partial \varphi}{\partial x^k} & \text{(gravitational force)} \\ & & -m\delta_{ab}\mathrm{R}^a_k\ddot{\varepsilon}^b & \text{(translational inertial force)} \\ & & +m[\vec{\omega} \wedge (\vec{\omega} \wedge \vec{x})]_k & \text{(centrifugal force)} \\ & & +m[\dot{\vec{\omega}} \wedge \vec{x}]_k \quad . & \text{(Jacobi force)} \end{array} \tag{4.22}$$

Finally, the equations of motion, which are the transformed of Newton's equations (4.6) in this generalized reference frame, are:

$$m\ddot{x}^k = -m\delta^{kl}\frac{\partial\varphi}{\partial x^l} - m\delta^{kl}\mathrm{R}^a_l\ddot{\varepsilon}^a + m[\vec{\omega}\wedge(\vec{\omega}\wedge\vec{x})]^k + 2m[\vec{\omega}\wedge\dot{\vec{x}}]^k + m[\dot{\vec{\omega}}\wedge\vec{x}]^k \quad . \tag{4.23}$$

On the other hand the Kinematical group of the subclass of *rigid observers*, called *Galilean* by Wheeler, is defined by Eqs.(4.16). Finally, the Kinematical group of *Galilean observers* (Wheeler's *absolute Galilean*) is defined by $R^a_i = \mathrm{const}$ and $\varepsilon^a = \mathrm{const}$.

When $\dot{\mathrm{R}}^a_i = 0$, the modification of Eq.(4.6) is due only to the *translational-inertial force*. Of course, corresponding to a transformation of the form (4.9), metric, affinity, and *dreibeins* are equivalent to the global flat ones and therefore do not represent true additional dynamical variables. It will be interesting to see whether it is possible to build up a Galilean theory in which the metric and the fields A_i assume an intrinsic dynamical content. This question will be dealt with in the following Sections.

5 Galilean limit of the relativistic mass-point theory

In order to understand the generality of the results so far obtained, it is profitable to make recourse to the axiomatic formulation of the so-called *Newtonian space-time structures* and to reconsider our formulation as a suitable non-relativistic limit of a Poincaré invariant theory. Axiomatic formulations of the possible geometries of Newtonian space-times has been introduced by Havas [2], Trautman [4],Künzle [5],and Kuchař [6].

Following Künzle, we define:

- A **Galilei structure** over a four-dimensional manifold $\mathcal{V}$ is a pair $(h^{\mu\nu}, t_\nu)$, where $h^{\mu\nu}$ is a symmetric covariant tensor of rank 3 and t_μ is a 1-form having the property that is the generator of the kernel of $h^{\mu\nu}$, $\forall x \in \mathcal{V}$.
  The triple $(\mathcal{V}; h^{\mu\nu}, t_\nu)$ is called a **Galilei Manifold**. A vector u^μ is called a *unit time-like vector* if $u^\mu t_\mu = 1$, and a contravariant tensor is called *space-like* if it vanish when contracted with t_μ on any index.

- A linear symmetric connection is called **a Galilei connection** ∇ if it is defined on a Galilei manifold $(\mathcal{V}; h^{\mu\nu}, t_\nu)$ and satisfies

$$\nabla_\rho h^{\mu\nu} = \nabla_\rho t_\mu = 0 \quad . \tag{5.1}$$

  It can be shown that such a connection exist if and only if t_μ is a closed 1-form and it is uniquely defined up to an arbitrary two-form χ according to:

$$\Gamma^\alpha_{\beta\gamma} = \overset{u}{\Gamma^\alpha_{\beta\gamma}} + t_\beta h^{\alpha\rho}\chi_{\rho\gamma} + t_\gamma h^{\alpha\rho}\chi_{\rho\beta} \quad , \tag{5.2}$$

  where $\Gamma^\alpha_{\beta\gamma}$ are the connection coefficients and

$$\overset{u}{\Gamma^\alpha_{\beta\gamma}} \equiv h^{\alpha\rho}\left[\frac{1}{2}h_{\beta\mu}h_{\gamma\nu}h^{\mu\nu}_{,\rho} + u^\sigma_{,\sigma}h_{\rho(\beta}t_{\gamma)}\right] - h_{\rho(\beta}h^{\alpha\rho}_{,\gamma)} - t_\beta u^\alpha_{,\gamma)} \ ; \tag{5.3}$$

  here u^α $(u^\alpha t_\alpha = 1)$ is an arbitrary given time-like unit vector field, and $\gamma_{\alpha\beta}$ is the associated covariant space metric defined by

$$\gamma_{\alpha\beta}u^\beta = 0 \qquad \text{and} \qquad \gamma_{\alpha\rho}\gamma^{\rho\beta} = \delta^\beta_\alpha - t_\alpha u^\beta \quad . \tag{5.4}$$

- A symmetric Galilei connection ∇ is called **Newtonian**, and the quadruple $(\mathcal{V}; h^{\mu\nu}, t_\nu; \Gamma)$ is called correspondingly a **Newtonian Manifold**, if the two-form χ in Eq.(5.2) is closed.

- Newtonian manifolds in which t_μ is also *exact* will be called *Special Newtonian Manifolds*: the hypothesis that t_μ be an *exact* one form implies the existence of a *global absolute time*. The standard Newtonian connection is indeed obtained by choosing $\chi_{\alpha\beta} = \varphi_{,\alpha}t_\beta - \varphi_{,\beta}t_\alpha$ where φ is the Newton's potential.

- Finally, the covariant *space metric* can be introduced by means of the relations:

$$\begin{aligned} h_{\mu\nu}h^{\nu\rho} &= \delta^\rho_\mu - t_\mu u^\rho \\ h_{\mu\nu}u^\nu &= 0 \quad . \end{aligned} \tag{5.5}$$

It is clear that the contravariant metric allows to assign lengths to *space-like* vectors but no lengths whatsoever to *time-like* vectors. These being the premises, we have:

1) the field equations, i.e. Newton's law of gravitation $\Delta\varphi = 4\pi G\rho$ (ρ mass density), can be rewritten as $R_{\alpha\beta} = 4\pi G\rho t_\alpha t_\beta$;

2) the matter Lagrangian for a mass-point, in the given Newtonian space-time, is (see Ref. [6]):

$$\begin{array}{rcl} A_M &=& \int d\lambda L_M(\lambda) \quad , \\ L_M &=& \dfrac{m}{2}\left(\dfrac{1}{t_\rho x'^\rho}\right) h_{\mu\nu} x'^\mu x'^\nu - m\varphi\left(t_\rho x'^\rho\right) \quad . \end{array} \tag{5.6}$$

Note that only within *Special Newtonian Manifolds* the first of Eqs.(5.6) can be rewritten in the form $A_M = \int dT \tilde{L}_M(T)$.

At this point, if our fields g_{ij}, Θ, A_i, A_0 are supposed to be resident within a *Special Newtonian* space-time, it is easy to make all the identifications which are relevant to our formulation. In particular, in a convenient coordinate chart $x^\mu = [t; x^k]$ of the *Special Newtonian* space time, we have:

$$\left\{ \begin{array}{rclrcl} \varphi &=& -\dfrac{1}{\Theta^2}\left(A_0 - \dfrac{1}{2}g^{ij}A_iA_j\right) & & & \\ t_\mu &=& [\Theta(t); \vec{0}] = [\dot{T}; \vec{0}] = t_{,\mu} & u^\mu &=& \dfrac{1}{\Theta}[1; -g^{ij}A_j] \\ h^{\mu\nu} &=& \begin{vmatrix} 0 & 0 \\ 0 & g^{ij} \end{vmatrix} & h_{\mu\nu} &=& \begin{vmatrix} g^{rs}A_rA_s & A_j \\ A_i & g_{ij} \end{vmatrix} \\ [\nabla_0]^i_0 &=& g^{ij}\left[\dfrac{\partial A_j}{\partial t} - \dfrac{1}{\Theta}\dfrac{\partial \Theta}{\partial t}A_j - \dfrac{\partial A_0}{\partial x^j}\right] & [\nabla_0]^0_0 &=& \dfrac{1}{\Theta}\dfrac{d\Theta}{dt} \\ [\nabla_0]^i_k &=& \dfrac{1}{2}g^{ij}\left[A_{j,k} - A_{k,j} + \dfrac{\partial g_{jk}}{\partial t}\right] & [\nabla_0]^0_k &=& 0 \\ [\nabla_j]^i_k &=& \Gamma^i_{jk} & [\nabla_j]^0_k &=& 0 \quad . \end{array} \right. \tag{5.7}$$

so that the Lagrangian (5.6) becomes:

$$\begin{array}{rcl} L^g_M &=& \dfrac{1}{t_\rho x'^\rho}\dfrac{m}{2}h_{\mu\nu}x'^\mu x'^\nu - m\varphi t_\rho x'^\rho \\ &=& \dfrac{m}{\Theta t'}\left[\dfrac{1}{2}g_{ij}x'^i x'^j + A_i x'^i t' + A_0 t' t'\right] \quad , \end{array} \tag{5.8}$$

which is precisely our Lagrangian (3.33) (see also Eqs.(4.12)).

We are now in a position to perform (see for example [14]) in a general way the non-relativistic limit of the mass-point Lagrangian that is invariant with respect to the Poincaré group *gauged á la Utiyama*. We have:

$$L^{Rg}_M = -mc\sqrt{-g_{\mu\nu}x'^\mu x'^\nu} \quad , \tag{5.9}$$

where we we have assumed the $(-1, +1, +1, +1)$ convention for the signature of the metric $g_{\mu\nu}$.

In order to perform the limit, we must write the explicit dependence of the metric tensor on c^2 (c velocity of light). As shown by Dautcourt [7],Ehlers [8] and Künzle [17], the correct parameterization to start with is:

$$g_{\mu\nu} = -c^2 t_\mu t_\nu + \overset{\circ}{g}_{\mu\nu} \quad , \tag{5.10}$$

and, for the inverse metric,

$$g^{\mu\nu} = h^{\mu\nu} - \frac{1}{c^2}\kappa^{\mu\nu} \quad . \tag{5.11}$$

From the relation

$$g^{\mu\nu} g_{\nu\rho} = \delta^\mu_\rho \quad , \tag{5.12}$$

we see that, at first order in $1/c^2$, the following identities are fulfilled:

$$\left\{ \begin{array}{lcl} h^{\mu\nu} t_\nu & = & 0 \\ h^{\mu\nu} \overset{\circ}{g}_{\nu\rho} & = & \delta^\mu_\rho - t_\rho \kappa^{\mu\nu} t_\nu \quad . \end{array} \right. \tag{5.13}$$

Therefore, in the limit $c \rightarrow +\infty$, we must identify $h^{\mu\nu}$ with the Newtonian *space metric* and $t_\mu t_\nu$ with the Newtonian *time metric*.

All the objects of the standard (*Special Newtonian Manifold* $t_\mu = t_{,\mu}$) Newtonian theory can then be reconstructed by the relations:

$$\left\{ \begin{array}{rcl} u^\mu t_{,\mu} & = & 1 \\ u^\nu \overset{0}{g}_{\nu\rho} & = & 2\, \varphi\, t_{,\rho} \\ h^{\mu\nu} \overset{0}{g}_{\nu\rho} & = & \delta^\mu_\rho - t_{,\rho}\, \kappa^{\mu\nu}\, t_{,\nu} \\ h_{\mu\nu} & = & \overset{\circ}{g}_{\mu\nu} - 2\varphi\, t_{,\mu} t_{,\nu} \end{array} \right. \tag{5.14}$$

that give the expressions $\varphi, u^\mu, h^{\mu\nu}, h_{\mu\nu}$ as functions of $t_{,\mu}$ and $\overset{\circ}{g}_{\mu\nu}$. The fields so defined automatically fulfill all the conditions required for the underlying space-time to be a **Newtonian manifold**. Within the coordinate chart used in Eqs.(5.7), it results

$$\overset{\circ}{g}_{\mu\nu} = h_{\mu\nu} + \begin{vmatrix} 2\varphi\Theta^2 & 0 \\ 0 & 0 \end{vmatrix} \quad . \tag{5.15}$$

By inserting the metric tensor (5.10) into the relativistic Lagrangian (5.9) and taking into account Eqs.(5.14), the expansion in terms of c^2 becomes:

$$L = c^2 \left[-m t_{,\nu}\, x'^\nu \right] + \left[\frac{1}{t_{,\rho} x'^\rho} \frac{m}{2} h_{\mu\nu} x'^\mu x'^\nu - m\, \varphi\, t_{,\rho} x'^\rho \right] + O(1/c^2) \quad . \tag{5.16}$$

We see that the zero$\underline{\text{th}}$ order part in c^2, reproduces the matter Lagrangian (5.6), while the coefficient of c^2 (which in the adapted frame (5.7) becomes $\left[-m\frac{dT}{dt}\frac{dt}{d\lambda} = -m\Theta\frac{dt}{d\lambda}\right]$) is, within the standard picture, the *central charge* $-m\Theta$ of the extended Galilei Group ($-m$ if one chooses $t = T$).

6 Searching for field equations

The expressions of the fields appearing in Eq.(4.12) are the generic-frame expressions of external fields resident in a spatially flat Newtonian space. On the other hand, having in mind the relations among the three-dimensional *gauge* fields that we have introduced and the four-dimensional metric, as contained in Eqs.(5.7,5.10,5.14), it seems natural to look for dynamical field equations in three dimensions by exploiting some limiting procedure over a four-dimensional theory. Now, let us observe that the contraction $c \to \infty$ from the Poincaré algebra to the extended Galilei algebra is a well defined procedure in the case of the single mass-point that we have studied in the previous section, due to the fact that the action is a Poincaré invariant: it amounts indeed to a uniquely defined contraction on the scalar representation. On the other hand, the field equations of motion do not transform like a scalar representation and it is well known that contracting a non-trivial representation is a delicate matter: a-priori, different contractions of the same equations could result. Therefore taking into account the fact that the Galilean matter Lagrangian (5.6) is nothing but the zero$^{\text{th}}$ order term of the $1/c^2$ expansion (5.16) of the general relativistic mass-point Lagrangian, we will build up the wanted Galilean variational problem by means of the zero$^{\text{th}}$-order term of the $1/c^2$ expansion of the full four-dimensional Einstein-Hilbert action for the gravitational field plus a single mass-point [18]:

$$\mathcal{S} = \mathcal{S}_F + \mathcal{S}_M = \frac{c^3}{16\pi G}\int d^4z\sqrt{-{}^4g}\;\; {}^4R - mc\int d\lambda\sqrt{-g_{\mu\nu}x'^{\mu}x'^{\nu}} \tag{6.1}$$

First of all, we shall restrict ourselves to globally hyperbolic space-time manifolds for which a *global* 3+1 splitting exists. The associated action principle will be formulated by re-expressing the Einstein-Hilbert Action (6.1) in the form given by Arnowitt-Deser-Misner

[19] and De-Witt [20]. This is tantamount to assume the existence of an absolute-time foliation of space-time ($t = t(z^\mu)$) and of a global coordinate system in which $t_{,\mu} = (\Theta(t), \vec{0})$ as in Eq.(5.6). In such a system, $g_{\mu\nu}$ must be of the form of Eq.(5.10) with $t_\mu = t_{,\mu}$ and the integration measure d^4z can be rewritten as dtd^3z.

Then, owing to Eqs.(5.7,5.10,5.14), we can put

$$\begin{aligned} g_{\mu\nu} &= -c^2 t_{,\mu} t_{,\nu} + \overset{\circ}{g}_{\mu\nu} \\ &= -c^2 \begin{vmatrix} \Theta^2 & 0 \\ 0 & 0 \end{vmatrix} + \begin{vmatrix} 2A_0 & A_j \\ A_i & g_{ij} \end{vmatrix} + \tfrac{1}{c^2} \begin{vmatrix} 2\alpha_0 & \alpha_j \\ \alpha_i & \gamma_{ij} \end{vmatrix} + \tfrac{1}{c^4} \begin{vmatrix} \dots & \dots \\ \dots & \beta_{ij} \end{vmatrix} + O(\tfrac{1}{c^6}) \end{aligned} \tag{6.2}$$

and

$$g^{\mu\nu} = \begin{vmatrix} 0 & 0 \\ 0 & g^{ij} \end{vmatrix} - \tfrac{1}{c^2\Theta^2} \begin{vmatrix} 1 & -g^{jk}A_k \\ -g^{ik}A_k & g^{ik}g^{jl}(\gamma_{kl} + A_kA_l) \end{vmatrix} - \tfrac{1}{c^4\Theta^4} \begin{vmatrix} A & \dots \\ \dots & \dots \end{vmatrix} + O(\tfrac{1}{c^6}) \ . \tag{6.3}$$

While some terms of order c^{-4} do contribute to the zero$^{\text{th}}$ order term we are interested in, *no term of order c^{-6} can survive the contraction*, so that they will be ignored from now on. Note that the parameterization (6.2) of $g_{\mu\nu}$ is the most general one which can be *locally* connected to the flat Minkowski metric $\eta_{\mu\nu}$ by means of a general coordinate transformation. By relaxing this last restriction, i.e. allowing for $\Theta = \Theta(t, \boldsymbol{z})$, one obtains a parameterization of $g_{\mu\nu}$, which needs also a Weyl transformation to be *locally* connected with $\eta_{\mu\nu}$. This corresponds in some way to allow for a classical analogue of the *dilaton* degree of freedom dynamically interacting with the other fields by paying the price of abandoning the Galilean interpretation of $\Theta(t)$ given in the previous section. Yet, this additional liberty allows for interesting results and will be exploited explicitly in the second paper of the present series.

Following Arnowitt, Deser and Misner [19],we write the covariant and the contravariant four-dimensional metric in the form:

$$\begin{aligned} {}^4g_{\mu\nu} &= \begin{vmatrix} -N^2 + {}^3g^{ij}N_iN_j & N_j \\ N_i & {}^3g_{ij} \end{vmatrix} \\ {}^4g^{\mu\nu} &= \begin{vmatrix} -\frac{1}{N^2} & \frac{{}^3g^{jk}N_k}{N^2} \\ \frac{{}^3g^{il}N^l}{N^2} & {}^3g^{ij} - \frac{{}^3g^{il}N_l {}^3g^{jk}N_k}{N^2} \end{vmatrix} \quad , \end{aligned} \tag{6.4}$$

where ${}^3g_{ij}{}^3g^{jk} = \delta_i^k$. Then, neglecting surface terms, the action (6.1) can be written:

$$\begin{aligned} \mathcal{S} &= \frac{c^3}{16\pi G}\int dt d^3z \sqrt{{}^3g} N \left[{}^3R + {}^3g^{ik}\ {}^3g^{jl}(K_{ij}K_{kl} - K_{ik}K_{jl})\right] \\ &\quad + mc\int d\lambda \sqrt{(N^2 - {}^3g^{ij}N_iN_j)t't' - N_i x'^i t' - {}^3g_{ij}x'^i x'^j} \quad , \end{aligned} \tag{6.5}$$

where:

$$\begin{cases} {}^3g &= \det {}^3g_{ij} \\ {}^3\Gamma^k_{ij} &= {}^3g^{kl}\dfrac{1}{2}({}^3g_{il,j} + {}^3g_{jl,i} - {}^3g_{ij,l}) \\ {}^3R_{ij} &= {}^3\Gamma^k_{ik,j} - {}^3\Gamma^k_{ij,k} + {}^3\Gamma^k_{il}{}^3\Gamma^l_{jk} - {}^3\Gamma^k_{ij}{}^3\Gamma^l_{lk} \\ {}^3R &= {}^3g^{ij}\ {}^3R_{ij} \\ K_{ij} &= \dfrac{1}{2N}\left({}^3\nabla_i N_j + {}^3\nabla_j N_i - \dfrac{\partial {}^3g_{ij}}{\partial t}\right) \quad (\textit{extrinsic curvature}) \ , \end{cases} \tag{6.6}$$

having denoted by ${}^3\nabla$ the covariant three-space derivative with respect to the Christoffel connection of ${}^3g_{ij}$ In terms of the notation we have previously introduced for the expansion of the covariant four-dimensional metric, we have

$$\begin{cases} {}^3g_{ij} &\equiv g_{ij} + \dfrac{1}{c^2}\gamma_{ij} + \dfrac{1}{c^4}\beta_{ij} + O(\dfrac{1}{c^6}) \\ {}^3R &\equiv R + \dfrac{1}{c^2}R_1(g_{ij},\gamma_{ij}) + \dfrac{1}{c^4}R_2(g_{ij},\gamma_{ij},\beta_{ij}) + O(\dfrac{1}{c^6}) \\ . \qquad N_i &\equiv A_i + \dfrac{1}{c^2}\alpha_i + \dfrac{1}{c^4}\beta_i + O(\dfrac{1}{c^6}) \\ N^2 &\equiv c^2\Theta^2 - 2A + \dfrac{2}{c^2}\left[\alpha_0 - g^{ij}\alpha_i A_j - \dfrac{1}{2}\gamma_{rs}g^{ri}g^{sj}A_iA_j\right] + O(\dfrac{1}{c^4}) \\ NK_{ij} &\equiv {}^3B_{ij} = B_{ij} + \dfrac{1}{c^2}B^{(1)}_{ij} + O(\dfrac{1}{c^4}) \quad , \end{cases} \tag{6.7}$$

where we have defined:

$$\begin{aligned} A &= A_0 - \tfrac{1}{2}g^{ij}A_iA_j \\ B_{ij} &= \tfrac{1}{2}[{}^3\nabla_i A_j + {}^3\nabla_j A_i - \tfrac{\partial g_{ij}}{\partial t}] \\ B^{(1)}_{ij} &= \tfrac{1}{2}[{}^3\nabla_i\alpha_j + {}^3\nabla_j\alpha_i - \tfrac{\partial\gamma_{ij}}{\partial t} - A_k g^{kl}({}^3\nabla_j\gamma_{il} + {}^3\nabla_i\gamma_{lj} - {}^3\nabla_l\gamma_{ij})] \ . \end{aligned} \tag{6.8}$$

Then, inserting eqs.(6.6-6.8) into eq.(6.5), it follows:

$$
\begin{aligned}
\mathcal{S} &= \mathcal{S}_F + \mathcal{S}_M = \\
&= c^4 \left\{ \frac{1}{16\pi G} \int dt d^3 z \sqrt{g} \Theta R \right\} \\
&\quad + c^2 \left\{ \frac{1}{16\pi G} \int dt d^3 z \sqrt{g} \left[\Theta R_1 + \frac{\Theta}{2} g^{ij} \gamma_{ij} R - \frac{A}{\Theta} R + \Theta g^{ik} g^{jl} (B_{ij} B_{kl} - B_{ik} B_{jl}) \right] \right. \\
&\qquad \left. - m \int d\lambda \Theta t' \right\} \\
&\quad + \left\{ \frac{1}{16\pi G} \int dt d^3 z \sqrt{g} \left[\Theta R_2 + \frac{1}{2} g^{ij} \gamma_{ij} \Theta R_1 + \frac{1}{2} g^{ij} \beta_{ij} \Theta R \right. \right. \\
&\qquad - \frac{1}{2} \Theta \left(g^{ik} g^{jl} - \frac{1}{2} g^{ij} g^{kl} \right) \gamma_{ij} \gamma_{kl} R - \frac{A}{\Theta} R_1 \\
&\qquad - \frac{1}{\Theta} \left(\alpha_0 - g^{ij} A_i \alpha_j + \frac{1}{2} A g^{ij} \gamma_{ij} + \frac{1}{2} \gamma_{ij} g^{il} g^{jm} A_l A_m \right) R - \frac{A^2}{2\Theta^3} R \\
&\qquad + \frac{2}{\Theta} g^{ik} g^{jl} (B_{ij} B^{(1)}_{kl} - B_{ik} B^{(1)}_{jl}) - \frac{2}{\Theta} g^{ik} g^{jr} \gamma_{rs} g^{sl} (B_{ij} B_{kl} - B_{ik} B_{jl}) \\
&\qquad \left. + \frac{A}{\Theta^3} g^{ik} g^{jl} (B_{ij} B_{kl} - B_{ik} B_{jl}) \right] \\
&\qquad \left. + m \int d\lambda \frac{m}{\Theta t'} \left[\frac{1}{2} g_{ij} (x'^i + g^{ik} A_k t')(x'^i + g^{ik} A_k t') + A_0 t' t' \right] \right\} \\
&\quad + O(1/c^2) \quad ,
\end{aligned}
\tag{6.9}
$$

so that the zero$^{\text{th}}$-order term, identified as the *total Action* $\tilde{\mathcal{S}}$, is:

$$
\begin{aligned}
\tilde{\mathcal{S}} &= \frac{1}{16\pi G} \int dt d^3 z \sqrt{g} \left[\Theta R_2 + \frac{1}{2} g^{ij} \gamma_{ij} \Theta R_1 + \frac{1}{2} g^{ij} \beta_{ij} \Theta R \right. \\
&\qquad - \frac{1}{2} \Theta \left(g^{ik} g^{jl} - \frac{1}{2} g^{ij} g^{kl} \right) \gamma_{ij} \gamma_{kl} R - \frac{A}{\Theta} R_1 \\
&\qquad - \frac{1}{\Theta} \left(\alpha_0 - g^{ij} A_i \alpha_j + \frac{1}{2} A g^{ij} \gamma_{ij} + \frac{1}{2} \gamma_{ij} g^{il} g^{jm} A_l A_m \right) R - \frac{A^2}{2\Theta^3} R \\
&\qquad + \frac{2}{\Theta} g^{ik} g^{jl} (B_{ij} B^{(1)}_{kl} - B_{ik} B^{(1)}_{jl}) - \frac{2}{\Theta} g^{ik} g^{jr} \gamma_{rs} g^{sl} (B_{ij} B_{kl} - B_{ik} B_{jl}) \\
&\qquad \left. + \frac{A}{\Theta^3} g^{ik} g^{jl} (B_{ij} B_{kl} - B_{ik} B_{jl}) \right] \\
&\quad + m \int d\lambda \frac{m}{\Theta t'} \left[\frac{1}{2} g_{ij} (x'^i + g^{ik} A_k t')(x'^i + g^{ik} A_k t') + A_0 t' t' \right] \quad .
\end{aligned}
\tag{6.10}
$$

We see that *27 fields survive the contraction*, namely Θ, A, A_i, g_{ij}, α_0, α_i, γ_{ij}, β_{ij}, where A has been used as independent variable instead of A_0; on the other hand β_0 and β_i disappear at this order.

Let us now look for the invariance of the Action. In section 3 we have shown that, provided we assume the transformation properties (3.43, the matter part of the Action (6.10) is invariant under the *gauged* Galilei transformation). Now, in order to formulate the *local Galilei invariance* for the action (6.10), appropriate *gauge* transformations for the fields γ_{ij}, β_{ij}, α_i and α_0 must be postulated. The correct choice is

$$\begin{aligned}
\delta\gamma_{ij} &= -\frac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}\gamma_{kj} - \frac{\partial\tilde{\eta}^k(\boldsymbol{x},t)}{\partial x^i}\gamma_{kj} \\
\delta\alpha_0 &= 2\dot{\varepsilon}\alpha_0 - \alpha_i\frac{\partial\tilde{\eta}^i}{\partial t} \\
\delta\alpha_i &= \dot{\varepsilon}\alpha_i - \alpha_j\frac{\partial\tilde{\eta}^j}{\partial x^i} - \gamma_{ij}\frac{\partial\tilde{\eta}^j}{\partial t} \quad ,
\end{aligned} \tag{6.11}$$

and, indeed, by direct calculation it turns out the the total variation of the Action (6.10) under the transformation δ defined by (3.7), (3.43) and (6.11) is given by:

$$\begin{aligned}
\delta\tilde{\mathcal{S}} &= \int dt d^3z \left\{ \dot{\varepsilon}\tilde{\mathcal{L}} + \Theta\mathcal{EL}_A\left(\frac{\partial\mathcal{F}}{\partial t} - A_r g^{rs}\frac{\partial\mathcal{F}}{\partial z^s}\right) + \Theta\mathcal{EL}_{A_i}\frac{\partial\mathcal{F}}{\partial z^i} \right. \\
&\left. + \frac{1}{8\pi G}\frac{\partial}{\partial z^i}\left(\frac{\sqrt{g}A}{\Theta^2}[B^{ij} - (\mathrm{Tr}B)g^{ij}]\frac{\partial\mathcal{F}}{\partial z^j}\right)\right\} \quad .
\end{aligned} \tag{6.12}$$

As a matter of fact, this result means that a *quasi-invariance* of the Galilean total Action (6.10holds, neglecting surface terms, *modulo* the Euler-Lagrange equations for the fields A_i ($\mathcal{EL}_{A_i}$) and A ($\mathcal{EL}_A$) that are given by:

$$\begin{aligned}
\mathcal{EL}_A &= \frac{\delta\tilde{\mathcal{S}}}{\delta A} \\
\mathcal{EL}_{A_i} &= \frac{\delta\tilde{\mathcal{S}}}{\delta A_i} - \frac{\partial}{\partial z^k}\frac{\delta\tilde{\mathcal{S}}}{\delta\partial_k A_i}
\end{aligned} \tag{6.13}$$

Let us remark that this peculiarity is precisely what it should be expected in the case of a variational principle corresponding to a *singular* Lagrangian.

By analogy to the free mass-point case, the terms $c^4 ... + c^2 ...$ of eqs.(6.9) could be rewritten as $c^2(\mathcal{M} + c^2\mathcal{N})$ where $\mathcal{M} + c^2\mathcal{N}$ ought to be interpreted as the central charge of the *asymptotic* Galilei group. To avoid an infinite central-charge, the theory should, in some sense, provide the condition $\mathcal{N} = 0$ automatically. The discussion of the *asymptotic* Galilei group will be dealt with in a separate paper. This analysis will require taking into account the $1/c^2$ expansion of the neglected surface terms, as they are needed in the evaluation of the asymptotic Poincaré group in the case of asymptotically-flat space-times (see for example ref.[23]).

7 Galilean Covariant Formulations of Newtonian Gravity

7.1 The Newtonian Theory in Arbitrary Reference Frames

In Section 4 it was shown that the field $\Theta(t)$ has no real dynamical content since its effect amounts only to a redefinition of the evolution parameter t in the expression $T(t) = \int_0^t d\tau\Theta(\tau)$. It is easy to show that this fact is still true for its role within the total Action (6.10). Indeed, if we redefine the fields A_0, A_i, α_0 and α_i as follows (see eq.(4.4); from now on ${}^3\nabla$ will be replace by ∇),

$$\left\{\begin{array}{rl} \tilde{A}_0 & \equiv \dfrac{A_0}{\Theta^2} \quad ; \qquad \tilde{A} \equiv \dfrac{A}{\Theta^2} \\ \tilde{A}_i & \equiv \dfrac{A_i}{\Theta} \\ \tilde{\alpha}_0 & \equiv \dfrac{\alpha_0}{\Theta^2} \quad ; \qquad \tilde{\alpha}_i \equiv \dfrac{\alpha_i}{\Theta} \\ \tilde{B}_{ij} & \equiv \dfrac{B_{ij}}{\Theta} = \dfrac{1}{2}\left[\nabla_i\tilde{A}_j + \nabla_j\tilde{A}_i - \dfrac{\partial g_{ij}}{\partial T}\right] \\ \tilde{B}^{(1)}_{ij} & \equiv \dfrac{B^{(1)}_{ij}}{\Theta} = \dfrac{1}{2}\left[\nabla_i\tilde{\alpha}_j + \nabla_j\tilde{\alpha}_i - \dfrac{\partial\gamma_{ij}}{\partial T} - \tilde{A}_k g^{kl}(\nabla_i\gamma_{jl} + \nabla_j\gamma_{il} - \nabla_l\gamma_{ij})\right] \quad , \end{array}\right. \tag{7.1}$$

the Action (6.10) becomes:

$$\begin{aligned} \tilde{\mathcal{S}} &= \textstyle\int dTd^3z\tilde{\mathcal{L}} \\ &= \frac{1}{16\pi G}\int dTd^3z\Big[\tilde{R}_2 - \tilde{A}\tilde{R}_1 \\ &\qquad - \sqrt{g}R\left(-\frac{\tilde{A}^2}{2} + \tilde{\alpha}_0 - g^{ij}\tilde{A}_i\tilde{\alpha}_j + \frac{1}{2}\tilde{A}g^{ij}\gamma_{ij} + \frac{1}{2}\gamma_{ij}g^{il}g^{jm}\tilde{A}_l\tilde{A}_m\right) \\ &\qquad + 2g^{ik}g^{jl}(\tilde{B}_{ij}\tilde{B}^{(1)}_{kl} - \tilde{B}_{ik}\tilde{B}^{(1)}_{jl}) - 2g^{ik}g^{jr}\gamma_{rs}g^{sl}(\tilde{B}_{ij}\tilde{B}_{kl} - \tilde{B}_{ik}\tilde{B}_{jl}) \\ &\qquad + \tilde{A}g^{ik}g^{jl}(\tilde{B}_{ij}\tilde{B}_{kl} - \tilde{B}_{ik}\tilde{B}_{jl})\Big] \\ &\quad + m\int dTd\boldsymbol{z}\ \left[\frac{1}{2}g_{ij}(\frac{dx^i}{dT} + g^{ik}\tilde{A}_k)(\frac{dx^i}{dT} + g^{ik}\tilde{A}_k) + \tilde{A}\right]\delta^3[\boldsymbol{z} - \boldsymbol{x}(T)] \quad , \end{aligned} \tag{7.2}$$

where, for future convenience, we have introduced the notations

$$\left\{\begin{array}{rcl} \tilde{R}_1(g,\gamma) & = & \sqrt{g}R_1(g,\gamma) + \dfrac{1}{2}g^{ij}\gamma_{ij}\sqrt{g}R \\ \tilde{R}_2(g,\gamma,\beta) & = & \sqrt{g}R_2(g,\gamma,\beta) + \dfrac{1}{2}g^{ij}\gamma_{ij}R_1 \\ & & +\dfrac{1}{2}\sqrt{g}R\left[g^{ij}\beta_{ij} - (g^{ik}g^{jl} - \dfrac{1}{2}g^{ij}g^{kl})\gamma_{ij}\gamma_{kl}\right] \quad . \end{array}\right. \tag{7.3}$$

It is then seen that the Action (7.2) is independent of $\Theta(t)$.

For future reference we give here the explicit expressions of the quantities $\tilde{R}_1$ and $\tilde{R}_2$. They are:

$$\left\{\begin{array}{rcl} \tilde{R}_1(g,\gamma) & = & \sqrt{g}\left[-\left(R^{ij}-\frac{1}{2}g^{ij}R\right)\gamma_{ij}+\left(g^{ik}g^{jl}-g^{ij}g^{kl}\right)\nabla_k\nabla_l\gamma_{ij}\right] \\ \tilde{R}_2(g,\gamma,\beta) & = & \sqrt{g}\left[-\left(R^{ij}-\frac{1}{2}g^{ij}R\right)\beta_{ij}+\left(g^{ik}g^{jl}-g^{ij}g^{kl}\right)\nabla_k\nabla_l\beta_{ij}\right] \\ & & +\frac{\sqrt{g}}{2}g^{ab}\gamma_{ab}\left[-\left(R^{ij}-\frac{1}{2}g^{ij}R\right)\gamma_{ij}+\left(g^{ik}g^{jl}-g^{ij}g^{kl}\right)\nabla_k\nabla_l\gamma_{ij}\right] \\ & & +\sqrt{g}g^{ab}\gamma_{ib}\gamma_{jb}\left(R^{ij}-\frac{1}{2}g^{ij}R\right) \\ & & +\sqrt{g}g^{ab}g^{ri}g^{js}\gamma_{rs}\left[\nabla_a\nabla_b\gamma_{ij}+\nabla_i\nabla_i\gamma_{ab}-\nabla_a\nabla_i\gamma_{jb}-\nabla_i\nabla_a\gamma_{bj}\right] \\ & & +\sqrt{g}g^{ab}g^{ij}g^{rs}\left[\nabla_r\gamma_{ij}\nabla_a\gamma_{sb}-\frac{1}{4}\nabla_r\gamma_{ij}\nabla_s\gamma_{ab}+\frac{3}{4}\nabla_r\gamma_{ia}\nabla_s\gamma_{jb}\right. \\ & & \qquad\qquad \left.-\frac{1}{2}\nabla_r\gamma_{ia}\nabla_j\gamma_{sb}-\nabla_r\gamma_{is}\nabla_a\gamma_{jb}\right] \quad ; \end{array}\right. \tag{7.4}$$

We shall deal now with the problem of investigating the true dynamical degrees of freedom of the theory by means of a constraint analysis within the Hamiltonian formalism. The canonical momenta $[\dot{f}=\frac{\partial f}{\partial T}]$ are defined by:

$$\left\{\begin{array}{rcl} \pi^{ij} & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{g}_{ij}} \\ & = & \frac{-\sqrt{g}}{16\pi G}\Big[(g^{ik}g^{jl}-g^{ij}g^{kl})-(g^{im}\gamma_{mn}g^{nk}g^{jl}+g^{ik}g^{jm}\gamma_{mn}g^{nl} \\ & & \qquad -g^{im}\gamma_{mn}g^{nj}g^{kl}-g^{ij}g^{km}\gamma_{mn}g^{nl})\Big]\tilde{B}^{(1)}_{kl} \\ & & +\frac{-\sqrt{g}\tilde{A}}{16\pi G}(g^{ik}g^{jl}-g^{ij}g^{kl})\tilde{B}_{kl} \\ \pi^{ij}_{\gamma} & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\gamma}_{ij}}=\frac{-\sqrt{g}}{16\pi G}(g^{ik}g^{jl}-g^{ij}g^{kl})\tilde{B}_{kl} \\ p_i & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{x}^i}=g_{ij}(\frac{dx^i}{dT}+g^{ik}\tilde{A}_k)\delta^3[\boldsymbol{z}-\boldsymbol{x}(T)] \quad , \end{array}\right. \tag{7.5}$$

and

$$\left\{\begin{array}{rclrclrcl} \pi_A & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\tilde{A}}}=0 & \pi_{\alpha_0} & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\tilde{\alpha}}_0}=0 & \pi^{ij}_{\beta} & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\beta}_{ij}}=0 \\ \pi^i & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\tilde{A}}_i}=0 & \pi^i_{\alpha} & = & \frac{\delta\tilde{\mathcal{S}}}{\delta\dot{\tilde{\alpha}}_i}=0 \quad . & & & \end{array}\right. \tag{7.6}$$

Since the Lagrangian $\tilde{\mathcal{L}}$ is independent of the corresponding *velocities*, the latter momenta define in fact 14 *primary* constraints.

The Dirac Hamiltonian density is given by:

$$\begin{aligned}\tilde{H}_d &= \frac{1}{16\pi G}\left\{\tilde{A}\tilde{R}_1 - \tilde{R}_2 + \sqrt{g}R\left(\frac{\tilde{A}^2}{2} + \tilde{\alpha}_0 - g^{ij}\tilde{A}_i\tilde{\alpha}_j + \frac{1}{2}\gamma_{ij}g^{ir}g^{js}\tilde{A}_r\tilde{A}_s\right)\right\} \\ &+\frac{32\pi G}{\sqrt{g}}\left(g_{ik}g_{jl} - \frac{1}{2}g_{ij}g_{kl}\right)\pi_\gamma^{ij}\pi^{kl} - \frac{16\pi G\tilde{A}}{\sqrt{g}}\left(g_{ik}g_{jl} - \frac{1}{2}g_{ij}g_{kl}\right)\pi_\gamma^{ij}\pi_\gamma^{kl} \\ &+\frac{32\pi G}{\sqrt{g}}\left(g_{ik}\gamma_{jl} - \frac{1}{2}g_{ij}\gamma_{kl}\right)\pi_\gamma^{ij}\pi_\gamma^{kl} \\ &+\left(\frac{1}{2m}g^{ij}p_ip_j - m\tilde{A}\right)\delta^3[\boldsymbol{z} - x(T)] \\ &-[\tilde{\alpha}_k - \tilde{A}_ig^{ij}\gamma_{jk}]\tilde{\Phi}^k - \tilde{A}_ig^{ij}\tilde{\phi}_k \\ &+\lambda^A\pi_A + \lambda^{\alpha_0}\pi_{\alpha_0} + \lambda_i\pi^i + \lambda_i^\alpha\pi_\alpha^i + \lambda_{ij}^\beta\pi_\beta^{ij} \quad ,\end{aligned} \tag{7.7}$$

where we have introduced *ad hoc* notations for the following important quantities

$$\left\{\begin{array}{lcl}\tilde{\phi}_k &=& 2g_{ij}\nabla_k\pi^{kl} + p_k\delta^3[\boldsymbol{z} - x(T)] + 2\nabla_r[\pi_\gamma^{rs}\gamma_{sk}] - \pi_\gamma^{rs}\nabla_k\gamma_{rs} \\ \tilde{\Phi}^k &=& \nabla_l\pi_\gamma^{kl} \quad .\end{array}\right. \tag{7.8}$$

We will apply now the Dirac-Bergmann procedure. By imposing time-conservation of the *primary* constraints, we obtain the 14 *secondary* (not all independent) ones:

$$\left\{\begin{array}{lcl}\dot{\pi}_{\alpha_0} &=& -\frac{1}{16\pi G}\sqrt{g}R \simeq 0 \\ \dot{\pi}_\beta^{ij} &=& = -\frac{1}{16\pi G}\left[R^{ij} - \frac{1}{2}g^{ij}R\right] \simeq 0 \quad . \\ \dot{\pi}_A &=& -\frac{\tilde{R}_1}{16\pi G} + m\delta^3[\boldsymbol{z} - x(T)] - \frac{\sqrt{g}\tilde{A}}{16\pi G}R + \frac{16\pi G}{\sqrt{g}}(g_{ik}g_{jl} - \frac{1}{2}g_{ij}g_{kl})\pi_\gamma^{ij}\pi_\gamma^{rs} \simeq 0 \\ \dot{\pi}^i &=& g^{ik}\tilde{\phi}_k + \frac{\sqrt{g}}{16\pi G}R\left(g^{ik}\tilde{\alpha}_k - g^{ik}\gamma_{kl}g^{lm}\tilde{A}_m\right) \simeq 0 \\ \dot{\pi}_\alpha^i &=& \tilde{\Phi}^k + \frac{\sqrt{g}}{16\pi G}Rg^{kl}\tilde{A}_l \simeq 0 \quad .\end{array}\right. \tag{7.9}$$

An equivalent, more expressive, set of 10 *secondary* constraints is:

$$\left\{\begin{array}{lcl}\chi_{\alpha_0} &\equiv& \sqrt{g}R \simeq 0 \\ \chi_R^{ij} &\equiv& \sqrt{g}R^{ij} - \frac{1}{3}\sqrt{g}R \simeq 0 \\ \chi_1 &\equiv& -\frac{1}{16\pi G}\tilde{R}_1 + m\delta^3[\boldsymbol{z} - x(T)] + \frac{16\pi G}{\sqrt{g}}(g_{ik}g_{jl} - \frac{1}{2}g_{ij}g_{kl})\pi_\gamma^{ij}\pi_\gamma^{rs} \simeq 0 \\ \tilde{\phi}_k &\simeq& 0 \\ \tilde{\Phi}^k &\simeq& 0 \quad .\end{array}\right. \tag{7.10}$$

By imposing time-conservation of the *secondary* constraints, we obtain the *tertiary* constraints in the form :

$$\left\{\begin{array}{lcl} \dot{\chi}_{\alpha_0} & \simeq & 0 \\ \dot{\chi}_R^{ij} & \simeq & -16\pi G\sqrt{g}\left[\frac{1}{2}\nabla^i\nabla^j\left(\frac{\pi_\gamma}{\sqrt{g}}\right)+\Delta\left(\frac{\pi_\gamma^{ij}}{\sqrt{g}}\right)-\frac{1}{2}g^{ij}\Delta\left(\frac{\pi_\gamma}{\sqrt{g}}\right)\right]\simeq 0 \\ \dot{\chi}_1 & \simeq & \sqrt{g}g^{kl}g^{rs}\nabla_k\nabla_l\gamma^{rs}-\frac{(16\pi G)^2}{2\sqrt{g}}g_{rs}\pi_\gamma^{rs}(g_{ik}g_{jl}-\frac{1}{2}g_{ij}g_{kl})\pi_\gamma^{ij}\pi_\gamma^{kl}\simeq 0 \\ \dot{\tilde{\phi}}_k & \simeq & 0 \\ \dot{\tilde{\Phi}}^k & \simeq & 0 \ , \end{array}\right. \tag{7.11}$$

where Δ is the three-dimensional Laplace operator ($\Delta = \nabla^k\nabla_k$). In this way we get seven (not independent) *tertiary* constraints. Note in this connection that, because of the constraint $R^{ij} \simeq 0$, the covariant derivatives commute on the constraint's surface. Also, recall finally that all the fields' momenta are tensor densities of weight $+1$.

At this point, it is expedient to introduced the well-known *transverse-traceless-decomposition* of symmetric tensors, which, due to the *secondary* constraint $\chi_R^{ij} \simeq 0$, can now be referred to the globally flat metric (on the constraints surface) g_{ij}. Specifically:

$$\left\{\begin{array}{lcl} \pi^{ij} & = & \pi_{TT}^{ij}+\frac{1}{2}\left[g^{ij}\pi_T-\Delta^{-1}\nabla^i\nabla^j\pi_T\right]+\nabla^i\pi_L^j+\nabla^j\pi_L^i \\ \gamma_{ij} & = & \gamma_{ij}^{TT}+\frac{1}{2}\left[g_{ij}\gamma^T-\Delta^{-1}\nabla_i\nabla_j\gamma^T\right]+\nabla_i\gamma_j^L+\nabla_j\gamma_i^L \\ \pi_\gamma^{ij} & = & \pi_{\gamma TT}^{ij}+\frac{1}{2}\left[g^{ij}\pi_{\gamma T}-\Delta^{-1}\nabla^i\nabla^j\pi_{\gamma T}\right]+\nabla^i\pi_{\gamma L}^j+\nabla^j\pi_{\gamma L}^i \\ \beta_{ij} & = & \beta_{ij}^{TT}+\frac{1}{2}\left[g_{ij}\beta^T-\Delta^{-1}\nabla_i\nabla_j\beta^T\right]+\nabla_i\beta_j^L+\nabla_j\beta_i^L \\ \pi_\beta^{ij} & = & \pi_{\beta TT}^{ij}+\frac{1}{2}\left[g^{ij}\pi_{\beta T}-\Delta^{-1}\nabla^i\nabla^j\pi_{\beta T}\right]+\nabla^i\pi_{\beta L}^j+\nabla^j\pi_{\beta L}^i \\ \lambda_{ij}^\beta & = & \lambda_{ij}^{\beta TT}+\frac{1}{2}\left[g_{ij}\lambda^{\beta T}-\Delta^{-1}\nabla_i\nabla_j\lambda^{\beta T}\right]+\nabla_i\lambda_j^{\beta L}+\nabla_j\lambda_i^{\beta L} \quad , \end{array}\right. \tag{7.12}$$

where $g^{ij}\gamma_{ij}^{TT} = \nabla^i\gamma_{ai}^{TT} = 0$, $g^{ij}\beta_{ij}^{TT} = \nabla^i\beta_{ai}^{TT} = 0$ and $g_{ij}\pi_{TT}^{ij} = \nabla_i\pi_{TT}^{ai} = 0$, $g_{ij}\pi_{\gamma TT}^{ij} = \nabla_i\pi_{\gamma TT}^{ai} = 0$, $g_{ij}\pi_{\beta TT}^{ij} = \nabla_i\pi_{\beta TT}^{ai} = 0$. In terms of these quantities, it can be seen that the chain that starts from the *primary* constraint $\pi_\beta^{ij} \simeq 0$ gets contributions only from the TT part, $\pi_{\beta TT}^{ij}$. Consistently, the *longitudinal* and *trace* parts, $\pi_{\beta L}^i$ and $\pi_{\beta T}$, do not generate any chain.

At this stage of the procedure, the constraints $\tilde{\Phi}^k \simeq 0$, $\dot{\chi}_R^{ij} \simeq 0$ can be rewritten as:

$$\left\{\begin{array}{lcl} \tilde{\Phi}^k & = & \nabla^k\nabla_i\pi_{\gamma L}^i+\Delta\pi_{\gamma L}^k \\ \dot{\chi}_R^{ij} & = & 16\pi G\left[\Delta\pi_{\gamma TT}^{ij}+\Delta\cdot(\nabla^i\pi_{\gamma L}^j+\nabla^j\pi_{\gamma L}^i)+g^{ij}\Delta\cdot\nabla_k\pi_{\gamma L}^k+\nabla^i\nabla^j\nabla_k\pi_{\gamma L}^k\right] \quad . \end{array}\right. \tag{7.13}$$

It is then apparent from these expressions that, provided that the asymptotic boundary conditions are such as to allow the inference $\Delta f = 0 \implies f = 0$, a condition that is also necessary to the inversion of the *transverse-traceless decomposition*, the constraints $\dot{\chi}^{ij}_R$ and $\tilde{\Phi}^k$ are equivalent to:

$$\left\{ \begin{array}{rcl} \pi^{ij}_{\gamma TT} & \simeq & 0 \quad , \\ \pi^i_{\gamma L} & \simeq & 0 \quad . \end{array} \right. \tag{7.14}$$

Therefore we have in fact only three independent *tertiary* constraints, namely $\dot{\chi}_1 \simeq 0$ and the two independent components of the first line of eqs.(7.14).

Using these conditions for reexpressing the constraints $\chi_1 \simeq 0$ and $\dot{\chi}_1 \simeq 0$ in terms of the *transverse traceless variables*, we obtain:

$$\left\{ \begin{array}{rcl} \chi_1 & = & \frac{\sqrt{g}}{16\pi G}\Delta\gamma^T + m\delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \\ & & + \frac{16\pi G}{\sqrt{g}}\frac{1}{4}\Big[(\pi_{\gamma T})^2 - (\Delta^{(-1)}\nabla^i\nabla^j\pi_{\gamma T}) \cdot (\Delta^{(-1)}\nabla_i\nabla_j\pi_{\gamma T})\Big] \simeq 0 \quad , \\ \dot{\chi}_1 & = & -\pi_{\gamma T} \cdot (\Delta \cdot \nabla^k\gamma^L_k) + \Big[\frac{1}{2}\nabla_i\nabla_j\gamma^T - (\nabla_i\nabla_j\nabla^k\gamma^L_k)\Big] \cdot (\Delta^{(-1)}\nabla^i\nabla^j\pi_{\gamma T}) \\ & & - \frac{16\pi G}{\sqrt{g}}\frac{\pi_{\gamma T}}{8}\Big[(\pi_{\gamma T})^2 - (\Delta^{(-1)}\nabla^i\nabla^j\pi_{\gamma T}) \cdot (\Delta^{(-1)}\nabla_i\nabla_j\pi_{\gamma T})\Big] \simeq 0 \quad . \end{array} \right. \tag{7.15}$$

At this stage we have the following situation: *i)* $\pi_{\beta T} \simeq 0$, $\pi^i_{\beta L} \simeq 0$ do not generate *secondary* constraints; *ii)* $\pi_{\alpha_0} \simeq 0$ generate the *secondary* $\chi_{\alpha_0} \simeq 0$ and no *tertiary*; *iii)* $\pi^i_\alpha \simeq 0$ generate the *secondaries* $\tilde{\Phi}^i \simeq 0$, i.e. $\pi^i_{\gamma L} \simeq 0$ and no *tertiary*; *iv)* $\pi^i \simeq 0$ generate the *secondaries* $\tilde{\phi}_i \simeq 0$ and no *tertiary*; *v)* $\pi_A \simeq 0$ generates the *secondary* $\chi_1 \simeq 0$ and then the *tertiary* $\dot{\chi}_1 \simeq 0$; *vi)* $\pi^{ij}_{\beta TT} \simeq 0$ (only two independent constraints) generate $\chi^{ij}_R \simeq 0$ (only two independent constraints due to the Bianchi identities) and then the two *tertiaries* $\dot{\chi}^{ij}_R \simeq 0$.

We have only to find the *quaternary* constraints generated by the time derivatives of the *tertiary* constraints $\dot{\chi}_1 \simeq 0$ and $\dot{\chi}^{ij}_R \simeq 0$. While $\ddot{\chi}_1 \simeq 0$ is given in appendix A, we have:

$$\begin{array}{rcl} \ddot{\chi}^{ij}_R & = & \frac{\sqrt{g}}{2 \cdot 16\pi G}\,\Delta\Delta\gamma^{TT}_{ij} + \frac{16\pi G}{2\sqrt{g}}\,[\nabla^r\pi_{\gamma T}\nabla^s\pi_{\gamma T} \\ & & \quad - \nabla^r\nabla^s\nabla^i\nabla^j\Delta^{(-1)}\pi_{\gamma T}\nabla_i\nabla_j\Delta^{(-1)}\pi_{\gamma T} \\ & & \quad + \frac{1}{2}\nabla^i\nabla^r\Delta^{(-1)}\pi_{\gamma T}\nabla_i\nabla^s\pi_{\gamma T} + \frac{1}{2}\nabla^i\nabla^s\Delta^{(-1)}\pi_{\gamma T}\nabla_i\nabla^r\pi_{\gamma T} \\ & & \quad - \frac{1}{2}\nabla^r\nabla^s\Delta^{(-1)}\pi_{\gamma T}\Delta\pi_{\gamma T} + \frac{1}{2}\nabla^r\nabla^s\pi_{\gamma T}\Big] \simeq 0 \quad . \end{array} \tag{7.16}$$

Before ending the discussion of these chains of constraints, let us remark that the relevant sector of solutions of eqs.(7.15) is

$$\left\{\begin{array}{rcl} \pi_{\gamma_T} & \simeq & 0 \\ \sqrt{g}\Delta\gamma^T & \simeq & -16\pi G m \delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \quad . \end{array}\right. \tag{7.17}$$

Using eqs.(7.17) inside eqs.(7.16), we get $\ddot{\chi}^{ij}_R \simeq \frac{\sqrt{g}}{2\cdot 16\pi G}\, g^{ir}g^{js}\Delta\Delta\gamma^{TT}_{rs} \simeq 0$, which implies $\gamma^{TT}_{rs} \simeq 0$. By using eqs.(7.17) and $\gamma^{TT}_{rs} \simeq 0$ in $\ddot{\chi}_1$ (see Appendix A), we obtain

$$\begin{array}{rcl} \ddot{\chi}_1 & \simeq & \dfrac{\sqrt{g}}{16\pi G}\left[\left(\Delta\gamma^T - 2\Delta\cdot\nabla^k\gamma^L_k\right)g^{ij} + 2\nabla^i\nabla^j\nabla^k\gamma^L_k\right]\cdot \\ & & \cdot\nabla_i\nabla_j\left(\tilde{A}-\frac{1}{4}\gamma^T\right) \ , \end{array} \tag{7.18}$$

which implies $\tilde{A}-\frac{1}{4}\gamma^T \simeq 0$ as a relevant solution. Therefore we get in the end

$$\Delta\tilde{A} \simeq -4\pi G m\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \quad , \tag{7.19}$$

i.e. the Poisson equation in a *three-dimensional general covariant* form. This means that $\ddot{\chi}_1 \simeq 0$ is the equation which replaces the Poisson equation in an *arbitrary-absolute time respecting* frame; the important result just obtained is that, provided that the Newton potential $A_0 = -\varphi$, seen by the *Galilean* observers, is replaced by the effective potential $\tilde{A} = \frac{A_0}{\Theta^2} - \frac{1}{2}\frac{g^{ij}A_iA_j}{\Theta^2}$, then we get the Poisson equation for $\tilde{A}$ as the most relevant solution (see Eq. 4.15) in every *allowed* reference frame.

Some words should be spent about the “invariance” of Eq.(7.19). Since we have shown in Eq.(6.12) that the Action is *quasi-invariant* modulo the equation of motion, one could expect that Eq.(7.19) be invariant under *all* the *local* Galilei transformations, just as all other equations are. Yet, this is not true because Eq.(6.12) is not invariant under *local* Galilei boosts because it gets contributions from the *cocycle* term. This does not invalidated the invariance of the theory, however. Indeed, since the Action is *quasi-invariant modulo* equations of motion, there is anyway a *conserved charge* associated to the boosts [22]. Therefore, the full invariance of Poisson equation should be accounted for by the transformations generated by these *conserved charge.*

Finally, time conservation of the *quaternary* constraints gives the *quinquenary* constraints. One of these latter, precisely that following from the χ_1 chain, fixes the *multiplier*

λ_A. On the other hand, the chain originated by χ^{rs}_R continues along three more time derivations. To avoid cumbersome expression, we give the simplified forms of the leading terms for the previous relevant sector; using all the constraints already worked out, it follows:

$$\left\{ \begin{array}{rcl} \overset{(3)}{\chi}_1 = \dfrac{d^3}{dT^3}\chi_1 & \simeq & \dfrac{\sqrt{g}}{16\pi G}\left[\left(\Delta\gamma^T - 2\Delta\cdot\nabla^k\gamma^L_k\right) g^{ij} + 2\nabla^i\nabla^j\nabla^k\gamma^L_k\right]\cdot\nabla_i\nabla_j\lambda_A + ... \simeq 0 \\ \overset{(3)}{\chi}_R{}^{rs} = \dfrac{d^3}{dT^3}\chi^{rs}_R & \simeq & \Delta\Delta\pi^{rs}_{TT} \simeq 0 \\ \overset{(4)}{\chi}_R{}^{rs} = \dfrac{d^4}{dT^4}\chi^{rs}_R & \simeq & \dfrac{\sqrt{g}g^{ri}g^{sj}}{16\pi G}\Delta\Delta\Delta\beta^{TT}_{ij} + ... \simeq 0 \\ \overset{(5)}{\chi}_R{}^{rs} = \dfrac{d^5}{dT^5}\chi^{rs}_R & \simeq & \dfrac{\sqrt{g}g^{ri}g^{sj}}{16\pi G}\Delta\Delta\Delta\lambda^{\beta TT}_{ij} + ... \simeq 0 \quad , \end{array}\right. \tag{7.20}$$

The last one ends the chain and fixes the *transverse-traceless* part $\lambda^{\beta TT}_{ij}$ of the *multipliers* λ^{ij}_{β}.

Therefore, since λ_A is determined from eq.(7.20, the chain of $\pi_A \simeq 0$ contains two pairs of *second class* constraints (π_A,$\ddot\chi_1$), (χ_1,$\dot\chi_1$)). On the other hand, each of the two independents chains of $\pi^{ij}_{\beta TT} \simeq 0$ contains three pairs of *second class* constraints ($\pi^{ij}_{\beta TT}$,$\overset{(4)}{\chi}_R{}^{rs}$), ($\chi^{rs}_R$,$\overset{(3)}{\chi}_R{}^{rs}$), ($\dot\chi^{rs}_R$,$\ddot\chi^{rs}_R$), since the sixth time derivative of these *primary* constraints determine the two independent Dirac multipliers $\lambda^{\beta TT}_{ij}$.

In conclusion there are 18 *first-class* constraints and 16 *second-class* constraints. While the variables $\tilde{A}$,γ^T,β^{TT}_{ij}, γ^{TT}_{ij} and two components of g_{ij} are determined by half of the *second-class* constraints (the other half determines their canonical momenta), the variables $\tilde{A}_i$, 3 of the g_{ij}, $\tilde{\alpha}_i$, γ^L_i, $\tilde{\alpha}_0$, π_T (conjugated to one of the g_{ij}), β^L_i, β^T are *gauge* variables (their conjugated variables are determined by the *first-class* constraints); correspondingly, the eleven Dirac multipliers $\lambda_i, \lambda^{\alpha_0}, \lambda^{\alpha}_i, \lambda^{\beta T}, \lambda^{\beta L}_i$ remain arbitrary.

Thus, apart from the *particle* degrees of freedom, no physical *field* degrees of freedom survive, as indeed it should be, and the role of the Newton potential is taken by $\tilde{A}$, which satisfies a Poisson equation in the most relevant sector of solutions. It would be interesting to see whether unconventional sectors are allowed corresponding to more general solutions for the gravitational potential[2].

The logical connections of the various constraints involved is described in Fig.1, which summarizes what is being fixed by each chains.

We can conclude this section by noting that, in force of eqs.(7.10) and (7.11), the condition for the finiteness of the central-charge term is indeed satisfied, and that $\mathcal{N} = 0$ holds. It remains an open task that of performing the $1/c^2$ expansion of the neglected surface term. It is likely that clarifying this issue will be relevant also to the understanding of the role of the *cocycle* contribution to the *local* Galilei transformations of the Poisson equation.

7.2 The Newtonian Theory in Galilean Reference Frames

Starting from the general scheme of the *27-fields* theory it is now interesting to see that, by confining to a post-Newtonian like [21] parameterization for the four-dimensional covariant metric tensor, defined by $\Theta = 1$, $g_{ij} = \delta_{ij}$, $A_i = 0$ and $A = -\varphi$ (i.e. the fields as seen by the *Galilean* observers: see Eq.(4.8)), one obtains the maximum of similarity to Newton's theory, i.e. a non-generally covariant formulation which is valid only in *Galilean* reference frame connected by *global* Galilei transformations. It should be clear, however, that in this way we are dealing in fact with a *different* variational problem with respect to the previous one. Putting

$$ {}^4g_{\mu\nu} = \begin{vmatrix} -c^2 - 2\varphi + \frac{2\alpha_0}{c^2} & \frac{\alpha_i}{c^2} \\ \frac{\alpha_i}{c^2} & \delta_{ij} + \frac{\gamma_{ij}}{c^2} + \frac{\beta_{ij}}{c^4} \end{vmatrix} \quad , \tag{7.21}$$

the explicit expressions of the quantities R_1 and R_2 defined in Eq.(6.7) become:

$$\left\{ \begin{array}{lll} R_1 & = & \delta^{ij}\delta^{rs}[\gamma_{ir,sj} - \gamma_{ij,rs}] \\ R_2 & = & \gamma_{lm}\delta^{lr}\delta^{ms}\delta^{ij}[\gamma_{ij,rs} + \gamma_{rs,ij} - 2\gamma_{ir,sj}] \\ & & +\delta^{lm}\delta^{rs}\delta^{ij}[\gamma_{rs,l}\gamma_{mi,j} - \frac{1}{4}\gamma_{ij,l}\gamma_{rs,m} - \gamma_{li,j}\gamma_{mr,s} \\ & & \qquad + \frac{3}{4}\gamma_{ir,l}\gamma_{js,m} - \frac{1}{2}\gamma_{ir,l}\gamma_{jm,s}] \\ & & +\delta^{ij}\delta^{rs}[\beta_{ir,sj} - \beta_{ij,rs}] \quad , \end{array} \right. \tag{7.22}$$

---

[2] This could possibly be of some interest in connection with the debate about the so-called fifth force


First class constraints
corresponding *gauge* variables
$\pi_{\beta T} \simeq 0$
END
β^T
$\pi^i_{\beta L} \simeq 0$
END
β^L_i
$\pi_{\alpha_0} \simeq 0$
$\sqrt{g}R \simeq 0$
END
$\tilde{\alpha}_0$ and π_T
$\pi^i_\alpha \simeq 0$
$\pi^i_{\gamma L} \simeq 0$
END
$\tilde{\alpha}_i$ and γ^L_i
$\pi^i \simeq 0$
$\tilde{\phi}_i \simeq 0$
END
$\tilde{A}_i$ and 3-g_{ij}
Second class constraints
$\pi_A \simeq 0$
$\chi_1 \simeq 0$
$\dot{\chi}_1 \simeq 0$
$\pi^{ij}_{\beta TT} \simeq 0$
$\chi_R \simeq 0$
$\dot{\chi}_R \simeq 0$
Multipliers Equations
$\ddot{\chi}_1 \simeq 0$
$\lambda_A \simeq ...$
$\ddot{\chi}^{rs}_R \simeq 0$
$\overset{(3)}{\chi}_R{}^{rs} \simeq 0$
$\overset{(4)}{\chi}_R{}^{rs} \simeq 0$
$\lambda^{\beta TT}_{ij} \simeq ..$

Figure 1: What is being fixed by the constraints' chains for the *27-fields* theory.

and the total action $\tilde{\mathcal{S}}$ (6.10) results:

$$\begin{aligned}
\tilde{\mathcal{S}} &\equiv \tfrac{1}{16\pi G}\int dtd^3z\mathcal{L}_f + m\int dtd^3z\mathcal{L}_m\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \\
&= \tfrac{1}{16\pi G}\int dtd^3z\left[(\varphi+\tfrac{1}{2}\delta^{ij}\gamma_{ij})R_1+R_2\right] \\
&\quad +m\int dtd^3z\left[\tfrac{1}{2}\delta_{ij}\dot{x}^i\dot{x}^j-\varphi\right]\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \quad .
\end{aligned} \tag{7.23}$$

It is seen that the matter Lagrangian $\mathcal{L}_m$ has precisely the form which is to be expected for a *Galilean observer* if Eq.(4.8) are inserted in Eq.(4.1). Therefore $(t, \boldsymbol{z})$ define a system of coordinates for a *Galilean* reference frame.

Note that:

(1) α_0 and α_i do not appear in the lagrangian $\mathcal{L}_f$;

(2) $\mathcal{L}_f$ depends on β_{ij} in a pure additive way through the term $\delta^{ij}\delta^{rs}[\beta_{ir,sj}-\beta_{ij,rs}]$ (see Eqs.(7.22)), which is again a *surface* term; moreover β_{ij} is not coupled to the other fields.

We can put accordingly $\alpha_0 = \alpha_i = \beta_{ij} = 0$ without altering the dynamics of this theory which indeed depends now on φ and γ_{ij} only.

Let us note, moreover, that the c^4-order term in the expansion (6.9) is automatically zero in this case, while the c^2-order term becomes:

$$\frac{1}{16\pi G}\int dtd^3z\left[R_1 - 16\pi Gm\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)]\right] \quad . \tag{7.24}$$

The resulting Lagrangian is:

$$\mathcal{L} = \frac{1}{16\pi G}\left[(\varphi+\frac{1}{2}\delta^{ij}\gamma_{ij})R_1+R_2'\right] + m\left[\frac{1}{2}\delta_{ij}\dot{x}^i\dot{x}^j-\varphi\right]\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \quad , \tag{7.25}$$

where $R_2' = R_2|_{\beta_{ij}=0}$, does not depend on the velocities $\dot{\varphi}$ and $\dot{\gamma}_{ij}$, so that all the field-momenta play the role of *primary* constraints.

The Hamiltonian formulation is defined by:

$$\left\{\begin{aligned}
\pi_\varphi &= 0 \\
\pi_\gamma^{rs} &= 0 \\
p_k &= m\delta_{kl}\dot{x}^l \\
H_c &= \tfrac{1}{16\pi G}\int d^3z\left[-(\varphi+\tfrac{1}{2}\delta^{ij}\gamma_{ij})R_1 - R_2'\right] \\
&\quad +\int d^3z\left[\tfrac{1}{2m}\delta^{ij}p_ip_j + m\varphi\right]\delta^3[\boldsymbol{z}-\boldsymbol{x}(t)] \quad ,
\end{aligned}\right. \tag{7.26}$$

so that we have the 7 *primary* constraints:

$$\left\{\begin{array}{lcl} \pi_\varphi & \simeq & 0 \\ \pi_\gamma^{rs} & \simeq & 0 \ . \end{array}\right. \tag{7.27}$$

The Dirac Hamiltonian is:

$$H_d = H_c + \int d^3z \left[\lambda^\varphi(\boldsymbol{z},t)\pi_\varphi + \lambda^\gamma_{rs}(\boldsymbol{z},t)\pi_\gamma^{rs}\right] \quad , \tag{7.28}$$

where the λ's are the Dirac multipliers. Time-conservation of these constraints generate the 7 *secondary* constraints:

$$\begin{array}{rcl} \chi_\varphi(\boldsymbol{z},t) & \equiv & \dot{\pi}_\varphi(\boldsymbol{z},t) = \{\pi_\varphi(\boldsymbol{z},t), H_d\} = \dfrac{1}{16\pi G} R_1 - m\delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \simeq 0 \\ \chi_\gamma^{ij}(\boldsymbol{z},t) & \equiv & \dot{\pi}_\gamma^{ij}(\boldsymbol{z},t) = \{\pi_\gamma^{ij}(\boldsymbol{z},t), H_d\} \\ & = & \dfrac{1}{16\pi G}\left\{\dfrac{1}{2}[\delta^{ir}\delta^{js} - \dfrac{1}{2}\delta^{ij}\delta^{rs}]\delta^{ab}[\gamma_{ab,rs} + \gamma_{rs,ab} - \gamma_{ar,sb} - \gamma_{as,rb}] \right. \\ & & \left. + [\delta^{ir}\delta^{js} - \delta^{ij}\delta^{rs}]\partial_r\partial_s\varphi\right\} \simeq 0 \quad , \end{array} \tag{7.29}$$

while their time conservation gives the following condition on λ^γ_{ij} and λ^φ:

$$\begin{array}{rcl} \psi_\varphi(\boldsymbol{z},t) & \equiv & \dot{\chi}_\varphi(\boldsymbol{z},t) = \{\chi_\varphi(\boldsymbol{z},t), H_d\} \\ & = & \frac{1}{16\pi G}\delta^{ij}\delta^{rs}[\lambda^\gamma_{ir,sj} - \lambda^\gamma_{ij,rs}] + p_k\delta^{kl}\partial_l\delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \simeq 0 \\ \psi_\gamma^{ij}(\boldsymbol{z},t) & \equiv & \dot{\chi}_\gamma^{ij}(\boldsymbol{z},t) = \{\chi_\gamma^{ij}(\boldsymbol{z},t), H_d\} \\ & = & \dfrac{1}{16\pi G}\left\{\dfrac{1}{2}[\delta^{ir}\delta^{js} - \dfrac{1}{2}\delta^{ij}\delta^{rs}]\delta^{ab}[\lambda^\gamma_{ab,rs} + \lambda^\gamma_{rs,ab} - \lambda^\gamma_{ar,sb} - \lambda^\gamma_{as,rb}] \right. \\ & & \left. + [\delta^{ir}\delta^{js} - \delta^{ij}\delta^{rs}]\partial_r\partial_s\lambda^\varphi\right\} \simeq 0 \quad . \end{array} \tag{7.30}$$

The *secondary* constraints are just the Euler-Lagrange field equations. Let us remark that the constraint $\chi_\varphi \simeq 0$ is just the expected condition for the vanishing of the c^2 term (7.24) (in the Lagrangian description this term vanishes because of the Euler-Lagrange equation for φ). On the other hand, the mass-point equations are

$$\dot{p}_k = \{p_k, H_d[T]\} = -m\partial_k\varphi \quad , \tag{7.31}$$

i.e. the standard Newton's equations with potential φ. Finally, evaluating R_1 from the contraction $\delta_{ij}\chi_\gamma^{ij} \simeq 0$ and by substituting it in the constraint $\chi_\varphi \simeq 0$, we obtain the classical Poisson equation for the potential $\varphi(\boldsymbol{z},t)$, i.e.:

$$\delta^{ij}\partial_i\partial_j\varphi(\boldsymbol{z},t) = 4\pi G m\delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \ . \tag{7.32}$$

Let us remark that, if we put an inhomogeneous solution of eq.(7.32) into eq.(7.31), one should get the motion of the particle under the usual field reaction; this gives rise to problems of self-energy similar to those of, e.g., the special relativistic electromagnetic case.

In spite of what could appear from Eqs.(7.30), not all the 7 *secondary* constraints are independent: as a matter of fact only four of them are independent and, correspondingly, only four *multipliers* are determined by eqs.(7.30). In order to see this explicitly, let us complete the constraint analysis of the theory. First of all, we note that three combination of the primary constraints, given by

$$\Pi^k = \partial_l \pi_\gamma^{kl} \ , \tag{7.33}$$

are *first class.* This can be easily checked thanks to the fact that the following six relations

$$\begin{array}{rcl} \partial_l \chi_\gamma^{kl} & \equiv & 0 \\ \partial_l \psi_\gamma^{kl} & \equiv & 0 \ . \end{array} \tag{7.34}$$

hold identically. Consequently, one has to expect that three of the γ_{ij} and three of the λ_{ij}^γ are free quantities. In order to evidentiate explicitly the multipliers and the fields that are determined by the constraints, it is profitable again to parameterize γ_{ij} and λ_{ij}^γ in terms of the *transverse-traceless decomposition* of symmetric tensors, as:

$$\begin{array}{rcl} \gamma_{ij} & = & \zeta_{ij}^{TT} + \frac{1}{2}\left[\delta_{ij}\zeta^T - \Delta^{-1}\zeta_{,ij}^T\right] + \zeta_{i,j} + \zeta_{j,i} \\ \lambda_{ij}^\gamma & = & \lambda_{ij}^{\gamma TT} + \frac{1}{2}\left[\delta_{ij}\lambda^{\gamma T} - \Delta^{-1}\lambda_{,ij}^{\gamma T}\right] + \lambda_{i,j}^\gamma + \lambda_{j,i}^\gamma \ , \end{array} \tag{7.35}$$

where $\delta^{ij}\zeta_{ij}^{TT} = \delta^{ij}\zeta_{ai,j}^{TT} = 0$ and $\delta^{ij}\lambda_{ij}^{\gamma TT} = \delta^{ij}\lambda_{ai,j}^{\gamma TT} = 0$. In terms of these quantities, the *secondary* and *tertiary* constraints (Eqs.(7.29) and (7.30)) become:

$$\left\{ \begin{array}{rcl} \chi_\varphi & \simeq & 0 \ \Rightarrow \ \Delta\zeta^T + 16\pi G m \delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \simeq 0 \\ \chi_\gamma^{ij} & \simeq & 0 \ \Rightarrow \ [\delta^{ir}\delta^{js} - \delta^{ij}\delta^{rs}]\partial_r\partial_s[\varphi - \frac{1}{4}\zeta^T] + \delta^{ir}\delta^{js}\delta^{lm}\zeta_{ij,lm}^{TT} \simeq 0 \ , \end{array} \right. \tag{7.36}$$

$$\left\{ \begin{array}{rcl} \psi_\varphi & \simeq & 0 \ \Rightarrow \ \Delta\lambda^{\gamma T} - 16\pi G p_k \delta^{kl}\partial_l \delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \simeq 0 \\ \psi_\gamma^{ij} & \simeq & 0 \ \Rightarrow \ [\delta^{ir}\delta^{js} - \delta^{ij}\delta^{rs}]\partial_r\partial_s[\lambda_\varphi - \frac{1}{4}\lambda^{\gamma T}] + \delta^{ir}\delta^{js}\delta^{lm}\lambda_{ij,lm}^{\gamma TT} \simeq 0 \ , \end{array} \right. \tag{7.37}$$

respectively. The *transverse-traceless* decomposition shows that the equations $\chi_\gamma^{ij} \simeq 0$ cannot be solved for the fields ζ_i and the multipliers λ_i^γ. In particular, as to the multipliers, we can solve only for:

$$\begin{aligned}
\lambda^{\varphi} &= \lambda^{\varphi}[z; p_k, x^k] \\
\lambda^{\gamma T} &= \lambda^{\gamma T}[z; p_k, x^k] \\
\lambda^{\gamma TT}_{ij} &= \lambda^{\gamma TT}_{ij}[z; p_k, x^k] \quad ,
\end{aligned} \tag{7.38}$$

where asymptotic boundary conditions for the λ's allowing for the inference $\Delta f = 0 \Longrightarrow f = 0$ have been assumed. Substituting these expressions for the multipliers, the Dirac Hamiltonian becomes:

$$\begin{aligned}
H_d = H_c + \int d^3z \Big[& \lambda^{\varphi}[z; p_k, x^k]\pi_{\varphi} + \lambda^{\gamma TT}_{ij}[z; p_k, x^k]\pi^{ij}_{\gamma} \\
& + \frac{1}{2}\left(\delta_{ij}\lambda^{\gamma T}[z; p_k, x^k] - \frac{1}{\nabla^2}\lambda^{\gamma T}_{,ij}[z; p_k, x^k]\right)\pi^{ij}_{\gamma} \\
& - 2\lambda^{\gamma}_i \partial_j \pi^{ij}_{\gamma} \Big] \quad ,
\end{aligned} \tag{7.39}$$

an expression which shows that the undetermined multipliers λ^{γ}_i are associated to the first class constraints Π^k, as it must be.

As a consequence, the variational problem must be independent of the quantities ζ_i of Eqs.(7.35). In fact, in terms of the *transverse-traceless* quantities, we have:

$$\left\{ \begin{aligned}
R_1 &= -\Delta\zeta^T \\
R_2' &= \frac{3}{8}\zeta^T\Delta\zeta^T - \frac{1}{4}\delta^{ij}\delta^{rs}\delta^{lm}\zeta^{TT}_{ir,l}\zeta^{TT}_{js,m} + \delta^{ij}\zeta_{i,j}\Delta\zeta^T + \frac{\partial F^k[\zeta^{TT}_{ij}, \zeta^T, \zeta_i]}{\partial z^k} \\
F^k &= \tfrac{3}{8}\delta^{ki}\zeta^T\zeta^T_{,i} - \tfrac{5}{16}\delta^{kl}\delta^{ij}\zeta^T_{,i}(\Delta^{-1}\zeta^T)_{,jl} + \tfrac{1}{16}\delta^{kl}\delta^{ij}\delta^{rs}(\Delta^{-1}\zeta^T)_{,ir}(\Delta^{-1}\zeta^T)_{,jsl} \\
&\quad - \tfrac{1}{2}\delta^{kl}\delta^{ij}\delta^{rs}\zeta^{TT}_{ir}\zeta^{TT}_{jl,s} + \delta^{kl}\delta^{ij}\delta^{rs}\zeta^{TT}_{ir}\zeta^{TT}_{js,l} - \tfrac{1}{4}\delta^{kl}\delta^{ij}\delta^{rs}(\Delta^{-1}\zeta^T)_{,ir}\zeta^{TT}_{jl,s} \\
&\quad - \tfrac{1}{2}\delta^{kl}\delta^{ij}\delta^{rs}(\Delta^{-1}\zeta^T)_{,i}\zeta^{TT}_{jl,rs} + \tfrac{1}{2}\delta^{kl}\delta^{ij}\zeta^T_{,i}\zeta^{TT}_{jl} + \delta^{kl}\delta^{ij}\zeta^T_{,l}\zeta_{i,j} \\
&\quad + \tfrac{1}{2}\delta^{kl}\delta^{ij}\zeta^T_{,i}\zeta_{l,j} + \tfrac{1}{2}\delta^{kl}\delta^{ij}\zeta^T_{,i}\zeta_{j,l} - \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{i,r}\zeta^{TT}_{jl,s} \\
&\quad - \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{r,i}\zeta^{TT}_{jl,s} + \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{i,r}\zeta^{TT}_{js,l} + \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{r,i}\zeta^{TT}_{js,l} \\
&\quad + \delta^{kl}\delta^{ij}\delta^{rs}\zeta^{TT}_{il,r}\zeta_{j,s} + \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{i,r}\zeta_{j,sl} - \delta^{kl}\delta^{ij}\delta^{rs}\zeta_{i,rs}\zeta_{j,l} \\
&\quad - \tfrac{1}{2}\delta^{kl}\delta^{ij}\delta^{rs}(\Delta^{-1}\zeta^T)_{,ir}\zeta_{l,js} + \tfrac{1}{2}\delta^{kl}\delta^{ij}\delta^{rs}(\Delta^{-1}\zeta^T)_{,il}\zeta_{r,js} \quad .
\end{aligned} \right. \tag{7.40}$$

Thus, neglecting the total divergence F^k, and thanks to suitable cancellations, the variational problem for a *Galilean observer*, can be reformulated, as an effective theory, only in terms of ζ^T and ζ^{TT}_{ij}, in the form:

$$\begin{aligned}
\mathcal{S} = & \tfrac{1}{16\pi G}\int dt d^3z \left[-\varphi\Delta\zeta^T - \tfrac{1}{8}\zeta^T\Delta\zeta^T - \tfrac{1}{4}\delta^{ij}\delta^{rs}\delta^{lm}\zeta^{TT}_{ir,l}\zeta^{TT}_{js,m}\right] \\
& + m\int dt d^3z \left[\tfrac{1}{2}\delta_{ij}\dot{x}^i\dot{x}^j - \varphi\right]\delta^3[\boldsymbol{z} - \boldsymbol{x}(t)] \quad .
\end{aligned} \tag{7.41}$$

Let us remark that in eq.(7.41) the term depending on ζ^{TT}_{ij} is decoupled from the other degrees of freedom. Therefore, in order to get a variational principle for the Poisson equation, only the *auxiliary*, non propagating, variable ζ^T is needed.

This theory turns out to be *quasi-invariant* under the global infinitesimal transformations, which constitute the kinematical group of the Galilean reference frames (2.3), as it should be, provided that φ is a scalar field and γ_{ij} is a covariant space 2-tensor, i.e.

$$\left\{ \begin{array}{rcl} \delta\varphi &=& 0 \\ \delta\gamma_{ij} &=& -\omega^l[{c_{li}}^k\gamma_{kj} + {c_{lj}}^k\gamma_{ik}] \quad . \end{array} \right. \tag{7.42}$$

As a consequence of these transformation properties, the *transverse traceless* components transform according to

$$\left\{ \begin{array}{rcl} \delta\zeta^T &=& 0 \\ \delta\zeta_i &=& -\omega^l {c_{li}}^k[\zeta_k - \frac{1}{2}(\Delta^{-1}\zeta^T)_{,k}] \\ \delta\zeta^{TT}_{ij} &=& -\omega^l[{c_{li}}^k\zeta^{TT}_{kj} + {c_{lj}}^k\zeta^{TT}_{ik}] \quad , \end{array} \right. \tag{7.43}$$

so that, finally,

$$\delta\mathcal{S} = m\int dt \frac{d}{dt}\left[\delta_{ij}v^i x^j\right] \quad . \tag{7.44}$$

8 Acknowledgments

Roberto De Pietri wishes to thank C. Rovelli, Al Janis and E.T. Newman for the hospitality kindly offered to him at the Department of Physics and Astronomy. Massimo Pauri would like to express his deep appreciation and thanks to the *Center for Philosophy of Science*, for the warm and stimulating intellectual atmosphere experienced there, and the generous partial support obtained during the completion of the present work at the University of Pittsburgh.

Appendix A: Explicit expression for the constraints $\ddot{\chi}_1$ of the 27-fields theory.

Using the notations: $f_{;i} = \nabla_i f$, $f^{;i} = \nabla^i f$ and $(K_\eta) = (16\pi G/\sqrt{g})$, the explicit expression of the constraints $\ddot{\chi}_1$ takes the form:

$$
\begin{aligned}
\ddot{\chi}_1 \simeq\ & -\frac{1}{(K_\eta)}\tilde{A}_{;i}{}^{i}\gamma^{T}{}_{;j}{}^{j}+\frac{1}{4\,(K_\eta)}\gamma^{T}{}_{;i}{}^{i}\gamma^{T}{}_{;j}{}^{j}+\\
& \frac{1}{(K_\eta)}\tilde{A}_{;ij}\gamma^{T}{}_{;}{}^{ij}-\frac{1}{4\,(K_\eta)}\gamma^{T}{}_{;ij}\gamma^{T}{}_{;}{}^{ij}-\\
& \frac{1}{2\,(K_\eta)}\gamma^{T}{}_{;ij}\gamma^{TTij}{}_{;k}{}^{k}-\frac{2}{(K_\eta)}\tilde{A}_{;i}{}^{i}\gamma^{L}{}_{j;}{}^{j}{}_{k}{}^{k}+\\
& \frac{1}{2\,(K_\eta)}\gamma^{T}{}_{;i}{}^{i}\gamma^{L}{}_{j;}{}^{j}{}_{k}{}^{k}+\frac{2}{(K_\eta)}\tilde{A}_{;ij}\gamma^{L}{}_{k;}{}^{ijk}-\\
& \frac{1}{2\,(K_\eta)}\gamma^{T}{}_{;ij}\gamma^{L}{}_{k;}{}^{ijk}-\frac{1}{(K_\eta)}\gamma^{TT}{}_{ij;k}{}^{k}\gamma^{L}{}_{l;}{}^{ijl}-\\
& \frac{2}{(K_\eta)}\gamma^{L}{}_{i;jk}{}^{k}\gamma^{L}{}_{l;}{}^{ijl}+\frac{2}{(K_\eta)}\gamma^{L}{}_{i;j}{}^{j}{}_{k}\gamma^{L}{}_{l;}{}^{ikl}+\\
& \frac{1}{2\,(K_\eta)}\gamma^{T}{}_{;ij}\gamma^{Li}{}_{;}{}^{j}{}_{k}{}^{k}-\frac{1}{2\,(K_\eta)}\gamma^{T}{}_{;ij}\gamma^{Lj}{}_{;}{}^{i}{}_{k}{}^{k}+\\
& \frac{(K_\eta)}{4}\tilde{A}_{;i}{}^{i}\Delta^{-1}\pi_{\gamma T;jk}\Delta^{-1}\pi_{\gamma T;}{}^{jk}-\frac{(K_\eta)}{16}\gamma^{T}{}_{;i}{}^{i}\Delta^{-1}\pi_{\gamma T;jk}\Delta^{-1}\pi_{\gamma T;}{}^{jk}+\\
& \frac{(K_\eta)}{2}\gamma^{T}{}_{;ij}\Delta^{-1}\pi_{\gamma T;}{}^{i}{}_{k}\Delta^{-1}\pi_{\gamma T;}{}^{jk}+\frac{(K_\eta)}{4}\gamma^{TT}{}_{ij;kl}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\Delta^{-1}\pi_{\gamma T;}{}^{kl}+\\
& \frac{(K_\eta)}{2}\gamma^{L}{}_{i;jkl}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\Delta^{-1}\pi_{\gamma T;}{}^{kl}-\frac{(K_\eta)}{8}\Delta^{-1}\gamma^{T}{}_{;ijkl}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\Delta^{-1}\pi_{\gamma T;}{}^{kl}+\\
& (K_\eta)\gamma^{L}{}_{i;}{}^{i}{}_{jk}\Delta^{-1}\pi_{\gamma T;}{}^{j}{}_{l}\Delta^{-1}\pi_{\gamma T;}{}^{kl}+\frac{(K_\eta)}{4}\gamma^{T}{}_{;i}\Delta^{-1}\pi_{\gamma T;jk}\Delta^{-1}\pi_{\gamma T;}{}^{ijk}+\\
& \frac{(K_\eta)}{2}\gamma^{TT}{}_{ij;k}\Delta^{-1}\pi_{\gamma T;}{}^{k}{}_{l}\Delta^{-1}\pi_{\gamma T;}{}^{ijl}+(K_\eta)\gamma^{L}{}_{i;jk}\Delta^{-1}\pi_{\gamma T;}{}^{k}{}_{l}\Delta^{-1}\pi_{\gamma T;}{}^{ijl}-\\
& \frac{(K_\eta)}{4}\Delta^{-1}\gamma^{T}{}_{;ijk}\Delta^{-1}\pi_{\gamma T;}{}^{k}{}_{l}\Delta^{-1}\pi_{\gamma T;}{}^{ijl}+\frac{(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}{}_{j}\Delta^{-1}\pi_{\gamma T;kl}\Delta^{-1}\pi_{\gamma T;}{}^{jkl}+\\
& \frac{(K_\eta)}{4}\gamma^{TT}{}_{ij}\Delta^{-1}\pi_{\gamma T;kl}\Delta^{-1}\pi_{\gamma T;}{}^{ijkl}+\frac{(K_\eta)}{2}\gamma^{L}{}_{i;j}\Delta^{-1}\pi_{\gamma T;kl}\Delta^{-1}\pi_{\gamma T;}{}^{ijkl}-\\
& \frac{(K_\eta)}{8}\Delta^{-1}\gamma^{T}{}_{;ij}\Delta^{-1}\pi_{\gamma T;kl}\Delta^{-1}\pi_{\gamma T;}{}^{ijkl}-\frac{(K_\eta)}{2}\Delta^{-1}\pi_{\gamma T;ij}\pi_{k}{}^{k}{}_{;}{}^{ij}+\\
& (K_\eta)\tilde{A}_{;ij}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\pi_{\gamma T}-\frac{5\,(K_\eta)}{8}\gamma^{T}{}_{;ij}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\pi_{\gamma T}-\\
& \frac{(K_\eta)}{2}\gamma^{TT}{}_{ij;k}{}^{k}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\pi_{\gamma T}-(K_\eta)\gamma^{L}{}_{i;jk}{}^{k}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\pi_{\gamma T}+\\
& \frac{(K_\eta)}{2}\gamma^{L}{}_{i;j}{}^{j}{}_{k}\Delta^{-1}\pi_{\gamma T;}{}^{ik}\pi_{\gamma T}-\frac{3\,(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}{}_{jk}\Delta^{-1}\pi_{\gamma T;}{}^{jk}\pi_{\gamma T}-\\
& \frac{(K_\eta)}{2}\gamma^{TT}{}_{ij;k}\Delta^{-1}\pi_{\gamma T;}{}^{ijk}\pi_{\gamma T}-(K_\eta)\gamma^{L}{}_{i;jk}\Delta^{-1}\pi_{\gamma T;}{}^{ijk}\pi_{\gamma T}+
\end{aligned}
$$

$$\frac{(K_\eta)}{4}\Delta^{-1}\gamma^{T}{}_{;ijk}\Delta^{-1}\pi_{\gamma T;}{}^{ijk}\pi_{\gamma T}+\frac{(K_\eta)}{2}\pi_{i}{}^{i}{}_{;j}{}^{j}\pi_{\gamma T}-\frac{5\,(K_\eta)}{4}\tilde{A}_{;i}{}^{i}\pi_{\gamma T}{}^{2}+$$

$$\frac{5\,(K_\eta)}{16}\gamma^{T}{}_{;i}{}^{i}\pi_{\gamma T}{}^{2}+\frac{(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}{}_{j}{}^{j}\pi_{\gamma T}{}^{2}+\frac{(K_\eta)^{3}}{8}\Delta^{-1}\pi_{\gamma T;ij}\Delta^{-1}\pi_{\gamma T;}{}^{ij}\pi_{\gamma T}{}^{2}-$$

$$\frac{(K_\eta)^{3}}{8}\pi_{\gamma T}{}^{4}-(K_\eta)\tilde{A}_{;i}\pi_{\gamma T}\pi_{\gamma T;}{}^{i}+(K_\eta)\tilde{A}_{;i}\Delta^{-1}\pi_{\gamma T;}{}^{i}{}_{j}\pi_{\gamma T;}{}^{j}-$$

$$\frac{(K_\eta)}{4}\gamma^{T}{}_{;i}\Delta^{-1}\pi_{\gamma T;}{}^{i}{}_{j}\pi_{\gamma T;}{}^{j}+\frac{(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}{}_{j}\pi_{\gamma T}\pi_{\gamma T;}{}^{j}-$$

$$(K_\eta)\gamma^{L}{}_{i;}{}^{i}{}_{j}\Delta^{-1}\pi_{\gamma T;}{}^{j}{}_{k}\pi_{\gamma T;}{}^{k}-\frac{(K_\eta)}{2}\tilde{A}\pi_{\gamma T}\pi_{\gamma T;i}{}^{i}+$$

$$\frac{(K_\eta)}{8}\gamma^{T}\pi_{\gamma T}\pi_{\gamma T;i}{}^{i}+\frac{(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}\pi_{\gamma T}\pi_{\gamma T;j}{}^{j}+$$

$$\frac{(K_\eta)}{2}\tilde{A}\Delta^{-1}\pi_{\gamma T;ij}\pi_{\gamma T;}{}^{ij}-\frac{(K_\eta)}{8}\gamma^{T}\Delta^{-1}\pi_{\gamma T;ij}\pi_{\gamma T;}{}^{ij}-$$

$$\frac{(K_\eta)}{4}\gamma^{TT}{}_{ij}\pi_{\gamma T}\pi_{\gamma T;}{}^{ij}-\frac{(K_\eta)}{2}\gamma^{L}{}_{i;j}\pi_{\gamma T}\pi_{\gamma T;}{}^{ij}+$$

$$\frac{(K_\eta)}{8}\Delta^{-1}\gamma^{T}{}_{;ij}\pi_{\gamma T}\pi_{\gamma T;}{}^{ij}-\frac{(K_\eta)}{2}\gamma^{L}{}_{i;}{}^{i}\Delta^{-1}\pi_{\gamma T;jk}\pi_{\gamma T;}{}^{jk}$$

References

[1] É. Cartan:“Sur les variétés à connexion affine et la théorie de la relativité generalisée (suite)”, Ann. Ècole Norm. Sup. 40,(1923),325-412

[2] P. Havas: “Four-Dimensional Formulations of Newtonian Mechanics and their Relation to the Special and the General Theory of Relativity”, Rev. Mod. Phys. **36**,(1964),938.

[3] J. L. Anderson: *Principles of Relativity Physics*, Academic Press, London 1967.

[4] A. Trautman, ”Theories of Space, Time and Gravitation” in *Lectures on General Relativity*, S. Deser and K.W. Ford, eds., Prentice-hall, Englewood Cliffs, 1965.

[5] H.P. Künzle: “Galilei and Lorentz Structures on Space-Time: Comparison of the Corresponding Geometry and Physics”, Ann. Inst. Henry Poincaré, **42**,(1972),337.

[6] K. Kuchař: “Gravitation, Geometry, and Non-Relativistic Quantum Theory”, Phys. Rev. ,**22D**,6,(1980),1285.

[7] G. Dautcourt: “Die Newtonske Gravitationstheorie als Strenger Grenzfall der Allgemeinen Relativitätheorie”, Acta Phys. Pol. **25**,5,(1964),637.

[8] J. Ehlers: “On Limit Relations between, and Approximative Explanations of, Physical Theories”, in *Logic, Methodology and Philosophy of Science*, VII, B. Marcus et al., eds., Elsiever, Amsterdam (1986),405.

[9] D.B. Malament: “Newtonian Gravity, Limits, and the Geometry of Space”, in *From Quarks to Quasars*, ed. R.G. Colodny, Pittsburgh University Press, Pittsburgh (1986)p.181.

[10] J.Earman and M.Friedman: ”The Meaning and Status of Newton’s Law of Inertia and the Nature of Gravitational Forces”, Philosophy of Science, **40** (1973), 329-359.

[11] R. Utiyama: “Invariant Theoretical Interpretation of Interactions”, Phys. Rev. **101**,(1956),1597.

[12] E.Kretschmann: “Über den Physikalischen Sinn der Relativitätspostulaten”, Ann. Phys. **53**,(1917),375.

[13] L. Michel: “Invariance in Quantum Mechanics and Group Extensions”, in *Group Theoretical Concepts and Methods in Elementary Particle Physics*, Gordon and Breach, New York (1964). See also: G.Marmo, G. Morandi, A. Simoni and E.C.G. Sudarshan: “ Quasi-Invariance and Central Extensions”, Phys. Rev. **D 37**,8,(1988),2196.

[14] A. Barducci, C. Casalbuoni and L. Lusanna: “Classical Spinning Particles Interacting with External Gravitational Field” , Nucl. Phys. **124 B**,(1977),521.

[15] C.W. Misner, K.S. Thorne and J.A. Wheeler: *Gravitation*, chapter 12, W.H. Freeman & Co., San Francisco 1973.

[16] T. Levi Civita: *The n-body problem in general relativity*, D. Reidel, Dordrecht (Holland) 1964.

[17] H.P. Künzle: “Covariant Newtonian Limit of Lorentz Space-Time”, Gen. Rel. and Grav., **7**,5,(1976),445.

[18] L.D. Landau and E.M. Liftschitz: *The Classical Theory of Fields*, Addison-Wesley, Cambridge (Mass.) 1951.

[19] R. Arnowitz, S. Deser, C.W. Misner: "The Dynamics of General Relativity" in *Gravitation: an Introduction to Current Research*, L. Witten, ed., John Wiley & Sons, New York 1962.

[20] B.S. De Witt: "Quantum Theory of Gravity. I. The Canonical Theory", Phys. Rev. **160**,5,(1967),1113.

[21] C.W. Misner, K.S. Thorne, and J.A. Wheeler: *Gravitation*, chapter 39, W.H. Freeman & Co., San Francisco 1973.

[22] L. Lusanna: "The Second Noether Theorem as the Basis of the Theory of Singular Lagrangians and Hamiltonians Constraints", Riv. Nuovo Cimento **14**,3,(1991),1.

[23] T. Regge and C. Teitelboim: "Role of Surface Integrals in the Hamiltonian Formulation of General Relativity", Annals of Physics **88**,(1974),286.

[24] M. Chaichian, D. Louis Martinez and L. Lusanna: "Dirac's Constrained System: The Classification of Second-Class Constraints", preprint Helsinki HU-TFT-93-5.

[25] R. Sugano, Y. Kagraoka and T. Kimura: "Gauge Transformations and Gauge-Fixing Condition in Constraint System", Int. J. of Mod. Phys. A **7**,(1992),61.